

**ANNEX I**  
**SUMMARY OF PRODUCT CHARACTERISTICS**

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

## 1. NAME OF THE MEDICINAL PRODUCT

Koselugo 10 mg hard capsules  
Koselugo 25 mg hard capsules

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

### Koselugo 10 mg hard capsules

Each hard capsule contains 10 mg of selumetinib (as hydrogen sulfate).

### Koselugo 25 mg hard capsules

Each hard capsule contains 25 mg of selumetinib (as hydrogen sulfate).

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Hard capsule.

### Koselugo 10 mg hard capsules

White to off-white, opaque, size 4 (approximately 14 mm x 5 mm), hard capsule, which has a centre band and is marked with “SEL 10” in black ink.

### Koselugo 25 mg hard capsules

Blue, opaque, size 4 (approximately 14 mm x 5 mm), hard capsule, which has a centre band and is marked with “SEL 25” in black ink.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Koselugo as monotherapy is indicated for the treatment of symptomatic, inoperable plexiform neurofibromas (PN) in paediatric patients with neurofibromatosis type 1 (NF1) aged 3 years and above.

### 4.2 Posology and method of administration

Treatment with Koselugo should be initiated by a physician experienced in the diagnosis and the treatment of patients with NF1 related tumours.

#### Posology

The recommended dose of Koselugo is 25 mg/m<sup>2</sup> of body surface area (BSA), taken orally twice daily (approximately every 12 hours).

Dosing is individualised based on BSA (mg/m<sup>2</sup>) and rounded to the nearest achievable 5 mg or 10 mg dose (up to a maximum single dose of 50 mg). Different strengths of Koselugo capsules can be combined to attain the desired dose (Table 1).

**Table 1. Recommended dose based on body surface area**

Body surface area (BSA) <sup>a</sup>	Recommended dose
0.55 – 0.69 m <sup>2</sup>	20 mg in the morning and 10 mg in the evening
0.70 – 0.89 m <sup>2</sup>	20 mg twice daily
0.90 – 1.09 m <sup>2</sup>	25 mg twice daily
1.10 – 1.29 m <sup>2</sup>	30 mg twice daily
1.30 – 1.49 m <sup>2</sup>	35 mg twice daily
1.50 – 1.69 m <sup>2</sup>	40 mg twice daily
1.70 – 1.89 m <sup>2</sup>	45 mg twice daily
≥ 1.90 m <sup>2</sup>	50 mg twice daily

<sup>a</sup> The recommended dose for patients with a BSA less than 0.55 m<sup>2</sup> has not been established.

Treatment with Koselugo should continue as long as clinical benefit is observed, or until PN progression or the development of unacceptable toxicity. There is limited data in patients older than 18, therefore continued treatment into adulthood should be based on benefits and risks to the individual patient as assessed by the physician. However, start of treatment with Koselugo in adults is not appropriate.

#### Missed dose

If a dose of Koselugo is missed, it should only be taken if it is more than 6 hours until the next scheduled dose.

#### Vomiting

If vomiting occurs after Koselugo is administered, an additional dose is not to be taken. The patient should continue with the next scheduled dose.

#### Dose adjustments

Interruption and/or dose reduction or permanent discontinuation of selumetinib may be required based on individual safety and tolerability (see sections 4.4 and 4.8). Recommended dose reductions are given in Table 2 and may require the daily dose to be divided into two administrations of different strength or for treatment to be given as a once daily dose.

**Table 2. Recommended dose reductions for adverse reactions**

Body surface area (BSA)	Initial Koselugo dose <sup>a</sup> (mg/twice daily)	First dose reduction (mg/dose)		Second dose reduction (mg/dose) <sup>b</sup>	
		Morning	Evening	Morning	Evening
0.55 – 0.69 m <sup>2</sup>	20 mg in the morning and 10 mg in the evening	10	10	10 mg once daily	
0.70 – 0.89 m <sup>2</sup>	20	20	10	10	10
0.90 – 1.09 m <sup>2</sup>	25	25	10	10	10
1.10 – 1.29 m <sup>2</sup>	30	25	20	20	10
1.30 – 1.49 m <sup>2</sup>	35	25	25	25	10
1.50 – 1.69 m <sup>2</sup>	40	30	30	25	20
1.70 – 1.89 m <sup>2</sup>	45	35	30	25	20
≥ 1.90 m <sup>2</sup>	50	35	35	25	25

<sup>a</sup> Based on BSA as shown in Table 1.

<sup>b</sup> Permanently discontinue treatment in patients unable to tolerate Koselugo after two dose reductions.

Dose modifications for the management of adverse reactions associated with this medicinal product are presented in Table 3.

**Table 3. Recommended dose modifications for adverse reactions**

<b>CTCAE Grade*</b>	<b>Recommended dose modification</b>
<b>Grade 1 or 2</b> (tolerable – can be managed with supportive care)	Continue treatment and monitor as clinically indicated
<b>Grade 2</b> (intolerable – cannot be managed with supportive care) or <b>Grade 3</b>	Interrupt treatment until toxicity is grade 0 or 1 and reduce by one dose level when resuming therapy (see Table 2)
<b>Grade 4</b>	Interrupt treatment until toxicity is grade 0 or 1, reduce by one dose level when resuming therapy (see Table 2). Consider discontinuation

\* Common Terminology Criteria for Adverse Events (CTCAE)

*Dose modification advice for left ventricular ejection fraction (LVEF) reduction*

In cases of asymptomatic LVEF reduction of  $\geq 10$  percentage points from baseline and below the institutional lower level of normal (LLN), selumetinib treatment should be interrupted until resolution. Once resolved, selumetinib should be reduced by one dose level when resuming therapy (see Table 2).

In patients who develop symptomatic LVEF reduction or a grade 3 or 4 LVEF reduction, selumetinib should be discontinued and a prompt cardiology referral should be carried out (see section 4.4).

*Dose modification advice for ocular toxicities*

Selumetinib treatment should be interrupted in patients diagnosed with retinal pigment epithelial detachment (RPED) or central serous retinopathy (CSR) with reduced visual acuity until resolution; reduce selumetinib by one dose level when resuming therapy (see Table 2). In patients diagnosed with RPED or CSR without reduced visual acuity, ophthalmic assessment should be conducted every 3 weeks until resolution. In patients who are diagnosed with retinal vein occlusion (RVO), treatment with selumetinib should be permanently discontinued (see section 4.4).

*Dose adjustments for co-administration with CYP3A4 or CYP2C19 inhibitors*

Concomitant use of strong or moderate CYP3A4 or CYP2C19 inhibitors is not recommended and alternative agents should be considered. If a strong or moderate CYP3A4 or CYP2C19 inhibitor must be co-administered, the recommended Koselugo dose reduction is as follows:

- If a patient is currently taking 25 mg/m<sup>2</sup> twice daily, dose reduce to 20 mg/m<sup>2</sup> twice daily.
- If a patient is currently taking 20 mg/m<sup>2</sup> twice daily, dose reduce to 15 mg/m<sup>2</sup> twice daily (see Table 4 and section 4.5).

**Table 4. Recommended dose to achieve 20 mg/m<sup>2</sup> or 15 mg/m<sup>2</sup> twice daily dose level**

Body surface area	20 mg/m <sup>2</sup> twice daily (mg/dose)		15 mg/m <sup>2</sup> twice daily (mg/dose)	
	Morning	Evening	Morning	Evening
0.55 – 0.69 m <sup>2</sup>	10	10	10 mg once daily	
0.70 – 0.89 m <sup>2</sup>	20	10	10	10
0.90 – 1.09 m <sup>2</sup>	20	20	20	10
1.10 – 1.29 m <sup>2</sup>	25	25	25	10
1.30 – 1.49 m <sup>2</sup>	30	25	25	20
1.50 – 1.69 m <sup>2</sup>	35	30	25	25
1.70 – 1.89 m <sup>2</sup>	35	35	30	25
≥ 1.90 m <sup>2</sup>	40	40	30	30

### Special populations

#### Renal impairment

Based on clinical trials no dose adjustment is recommended in patients with mild, moderate, severe renal impairment or those with end stage renal disease (ESRD) (see section 5.2).

#### Hepatic impairment

Based on clinical trials, no dose adjustment is recommended in patients with mild hepatic impairment. The starting dose should be reduced in patients with moderate hepatic impairment to 20 mg/m<sup>2</sup> BSA, twice daily (see Table 4). Koselugo is contraindicated for use in patients with severe hepatic impairment (see sections 4.3 and 5.2).

#### Ethnicity

Increased systemic exposure has been seen in adult Asian subjects, although there is considerable overlap with Western subjects when corrected for body weight. No specific adjustment to the starting dose is recommended for paediatric Asian patients, however these patients, should be closely monitored for adverse events (see section 5.2).

#### Paediatric population

The safety and efficacy of Koselugo in children less than 3 years of age has not been established. No data are available.

#### Method of administration

Koselugo is for oral use. It should be taken on an empty stomach with no food or drink other than water 2 hours prior to dosing and 1 hour after dosing (see sections 4.5 and 5.2).

The capsules should be swallowed whole with water. The capsules should not be chewed, dissolved, or opened, because this could impair drug release and affect the absorption of selumetinib.

Koselugo should not be administered to patients who are unable or unwilling to swallow the capsule whole. Patients should be assessed for their ability to swallow a capsule before starting treatment. Standard medicine swallowing techniques are expected to be sufficient to swallow selumetinib capsules. For patients who have difficulties swallowing the capsule, referral to an appropriate health care professional such as a speech and language therapist could be considered to identify suitable methods that can be tailored to the particular patient.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.  
Severe hepatic impairment (see sections 4.2 and 5.2).

#### 4.4 Special warnings and precautions for use

##### Left ventricular ejection fraction (LVEF) reduction

Asymptomatic decreases in ejection fraction have been reported in 26% of paediatric patients in the pivotal clinical trial. Median time to initial onset of these adverse reactions was 232 days. A small number of serious reports of LVEF reduction associated with selumetinib have been reported in paediatric patients who participated in an expanded access program (see section 4.8).

Paediatric patients with a history of impaired left ventricular function or a baseline LVEF below institutional LLN have not been studied. LVEF should be evaluated by echocardiogram before initiation of treatment to establish baseline values. Prior to starting selumetinib treatment, patients should have an ejection fraction above the institutional LLN.

LVEF should be evaluated at approximately 3-month intervals, or more frequently as clinically indicated, during treatment. Reduction in LVEF can be managed using treatment interruption, dose reduction or treatment discontinuation (see section 4.2).

##### Ocular toxicity

Patients should be advised to report any new visual disturbances. Adverse reactions of blurred vision have been reported in paediatric patients receiving selumetinib. Isolated cases of RPED, CSR and RVO in adult patients with multiple tumour types, receiving treatment with selumetinib monotherapy and in combination with other anti-cancer agents, and in a single paediatric patient with pilocytic astrocytoma on selumetinib monotherapy, have been observed (see section 4.8).

In line with clinical practice an ophthalmological evaluation prior to treatment initiation and at any time a patient reports new visual disturbances is recommended. In patients diagnosed with RPED or CSR without reduced visual acuity, ophthalmic assessment should be conducted every 3 weeks until resolution. If RPED or CSR is diagnosed and visual acuity is affected, selumetinib therapy should be interrupted and the dose reduced when treatment is resumed (see section 4.2). If RVO is diagnosed, treatment with selumetinib should be permanently discontinued (see section 4.2).

##### Liver laboratory abnormalities

Liver laboratory abnormalities, specifically AST and ALT elevations, can occur with selumetinib (see section 4.8). Liver laboratory values should be monitored before initiation of selumetinib and at least monthly during the first 6 months of treatment, and thereafter as clinically indicated. Liver laboratory abnormalities should be managed with dose interruption, reduction or treatment discontinuation (see Table 2 in section 4.2).

##### Skin and subcutaneous disorders

Skin rash (including maculopapular rash and acneiform rash), paronychia and hair changes have been reported very commonly in the pivotal clinical study (see section 4.8). Dry skin, hair colour changes, paronychia and rash maculo-papular were seen more frequently in younger children (age 3-11 years) and acneiform rash was seen more frequently in post-pubertal children (age 12-16 years).

##### Vitamin E supplementation

Patients should be advised not to take any supplemental vitamin E. Koselugo 10 mg capsules contain 32 mg vitamin E as the excipient, D-alpha-tocopheryl polyethylene glycol 1000 succinate (TPGS). Koselugo 25 mg capsules contain 36 mg vitamin E as TPGS. High doses of vitamin E may increase the risk of bleeding in patients taking concomitant anticoagulant or antiplatelet medicinal products (e.g., warfarin or acetylsalicylic acid). Anticoagulant assessments, including international normalised ratio or prothrombin time, should be conducted more frequently to detect when dose adjustments of the anticoagulant or antiplatelet medicinal products are warranted (see section 4.5).

#### Risk of choking

Selumetinib is available as a capsule which must be swallowed whole. Some patients, in particular children < 6 years of age, may be at risk of choking on a capsule formulation due to developmental, anatomical or psychological reasons. Therefore, selumetinib should not be administered to patients who are unable or unwilling to swallow the capsule whole (see section 4.2).

#### Women of child bearing potential

Koselugo is not recommended in women of child bearing potential who are not using contraception (see section 4.6).

### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in healthy adults (aged  $\geq$  18 years).

#### Active substances that may increase selumetinib plasma concentrations

Co-administration with a strong CYP3A4 inhibitor (200 mg itraconazole twice daily for 4 days) increased selumetinib  $C_{max}$  by 19% (90% CI 4, 35) and AUC by 49% (90% CI 40, 59) in healthy adult subjects.

Co-administration with a strong CYP2C19/moderate CYP3A4 inhibitor (200 mg fluconazole once daily for 4 days) increased selumetinib  $C_{max}$  by 26% (90% CI 10, 43) and AUC by 53% (90% CI 44, 63) in healthy adult subjects, respectively.

Concomitant use of erythromycin (moderate CYP3A4 inhibitor) or fluoxetine (strong CYP2C19/CYP2D6 inhibitor) is predicted to increase selumetinib AUC by ~30-40% and  $C_{max}$  by ~20%.

Co-administration with medicinal products that are strong inhibitors of CYP3A4 (e.g., clarithromycin, grapefruit juice, oral ketoconazole) or CYP2C19 (e.g., ticlopidine) should be avoided.

Co-administration with medicinal products that are moderate inhibitors of CYP3A4 (e.g., erythromycin and fluconazole) and CYP2C19 (e.g., omeprazole) should be avoided.

If co-administration is unavoidable, patients should be carefully monitored for adverse events and the selumetinib dose should be reduced (see section 4.2 and Table 4).

#### Active substances that may decrease selumetinib plasma concentrations

Co-administration with a strong CYP3A4 inducer (600 mg rifampicin daily for 8 days) decreased selumetinib  $C_{max}$  by -26% (90% CI -17, -34) and AUC by -51% (90% CI -47, -54).

Concomitant use of strong CYP3A4 inducers (e.g., phenytoin, rifampicin, carbamazepine, St. John's Wort) or moderate CYP3A4 inducers with Koselugo should be avoided.

#### Active substances whose plasma concentrations may be altered by selumetinib

*In vitro*, selumetinib is an inhibitor of OAT3. The potential for a clinically relevant effect on the pharmacokinetics of concomitantly administered substrates of OAT3 (e.g., methotrexate and furosemide) cannot be excluded (see section 5.2).

TPGS is a P-gp inhibitor *in vitro* and it cannot be excluded that it may cause clinically relevant drug interactions with substrates of P-gp (e.g., digoxin or fexofenadine).

The effect of selumetinib on the exposure of oral contraceptives has not been evaluated. Therefore, use of an additional barrier method should be recommended to women using hormonal contraceptives (see section 4.6).

#### Effect of gastric acid reducing agents on selumetinib

Selumetinib capsules do not exhibit pH dependent dissolution. Koselugo can be used concomitantly with gastric pH modifying agents (i.e., H<sub>2</sub>-receptor antagonists and proton pump inhibitors) without restrictions, except for omeprazole which is a CYP2C19 inhibitor.

#### Vitamin E

Koselugo capsules contain vitamin E as the excipient TPGS. Therefore, patients should avoid taking supplemental vitamin E and anticoagulant assessments should be performed more frequently in patients taking concomitant anticoagulant or antiplatelet medicinal products (see section 4.4).

### **4.6 Fertility, pregnancy and lactation**

#### Women of childbearing potential/Contraception in males and females

Women of childbearing potential should be advised to avoid becoming pregnant while receiving Koselugo. It is recommended that a pregnancy test should be performed on women of childbearing potential prior to initiating treatment.

Both male and female patients (of reproductive potential) should be advised to use effective contraception during and for at least 1 week after completion of treatment with Koselugo. It cannot be excluded that selumetinib may reduce the effectiveness of oral contraceptives, therefore women using hormonal contraceptives should be recommended to add a barrier method (see section 4.5).

#### Pregnancy

There are no data on the use of selumetinib in pregnant women. Studies in animals have shown reproductive toxicity including embryofetal death, structural defects and reduced foetal weights (see section 5.3). Koselugo is not recommended during pregnancy and in women of childbearing potential not using contraception (see section 4.4).

If a female patient or a female partner of a male patient receiving Koselugo becomes pregnant, she should be apprised of the potential risk to the foetus.

#### Breast-feeding

It is not known whether selumetinib, or its metabolites, are excreted in human milk. Selumetinib and its active metabolite are excreted in the milk of lactating mice (see section 5.3). A risk to the breast-fed child cannot be excluded, therefore breast-feeding should be discontinued during treatment with Koselugo.

#### Fertility

There are no data on the effect of Koselugo on human fertility. Selumetinib had no impact on fertility and mating performance in male and female mice, although a reduction in embryonic survival was observed in female mice (see section 5.3).

### **4.7 Effects on ability to drive and use machines**

Koselugo may have a minor influence on the ability to drive and use machines. Fatigue, asthenia and visual disturbances have been reported during treatment with selumetinib and patients who experience these symptoms should observe caution when driving or using machines.

### **4.8 Undesirable effects**

#### Summary of the safety profile

The safety profile of selumetinib monotherapy in paediatric patients with NF1 who have inoperable PN has been determined following evaluation of a combined safety population of 74 paediatric patients (20-30 mg/m<sup>2</sup> twice daily). This paediatric 'pool' of patients comprised 50 patients in SPRINT Phase II Stratum 1, treated with selumetinib 25 mg/m<sup>2</sup> twice daily (the pivotal dataset) and 24 patients in SPRINT Phase I treated with 20 to 30 mg/m<sup>2</sup> selumetinib twice daily (the dose finding



study). There were no clinically relevant differences in the safety profile between SPRINT Phase I and SPRINT Phase II Stratum 1. This safety profile was also substantiated by a pool of safety data from 7 AstraZeneca sponsored studies in adult patients with multiple tumour types (N = 347) who received 75 to 100 mg twice daily).

In the paediatric pool, the median total duration of selumetinib treatment in paediatric patients with NF1 who have PN was 55 months (range: < 1 to 97 months), 61% of patients were exposed to selumetinib treatment for > 48 months and 16% for >72 months. Patients aged  $\geq 2$  to 11 years (N = 45) had a higher incidence of the following adverse drug reactions (ADRs) compared to patients aged 12 to 18 years (N = 29): hypoalbuminaemia, dry skin, pyrexia, hair colour changes, rash maculopapular and paronychia.

In the paediatric pool (N = 74; comprising 50 patients from the pivotal SPRINT Phase II Stratum 1 dataset and 24 patients from the supportive SPRINT Phase I dataset), the most common adverse reactions of any grade (incidence  $\geq 45\%$ ) were vomiting (86%), diarrhoea (81%), blood creatine phosphokinase increased (77%), nausea (77%), dry skin (65%), pyrexia (61%), dermatitis acneiform (61%), asthenic events (59%), paronychia (57%), stomatitis (55%), haemoglobin decreased (54%), non-acneiform rashes (53%), hypoalbuminaemia (51%), and aspartate aminotransferase increased (51%). Dose interruptions and reductions due to adverse events were reported in 82% and 39% of patients, respectively. The most commonly reported ADRs leading to dose modification (dose interrupted or dose reduced) of selumetinib were vomiting (32%), paronychia (23%), nausea (19%), diarrhoea (15%) and pyrexia (11%). Permanent discontinuation due to adverse events was reported in 12% of the patients. The following serious adverse reactions were reported: diarrhoea (3%), anaemia (3%), pyrexia (3%), blood CPK increased (3%), blood creatinine increased (1%), oedema peripheral (1%) and vomiting (1%).

#### Tabulated list of adverse reactions

Table 5 presents the adverse reactions identified in the paediatric population with NF1 who have inoperable PN and in adult patients (see footnote to Table 5). The frequency is determined from the paediatric pool (N = 74); comprising 50 patients from the pivotal SPRINT Phase II Stratum 1 dataset and 24 patients from the supportive SPRINT Phase I dataset. Adverse drug reactions (ADRs) are organised by MedDRA system organ class (SOC). Within each SOC, preferred terms are arranged by decreasing frequency and then by decreasing seriousness. Frequencies of occurrence of adverse reactions are defined as: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ) and not known (cannot be estimated from available data), including isolated reports.

**Table 5. Adverse drug reactions reported in the paediatric pool (pivotal SPRINT Phase II Stratum 1 [N = 50] and supportive SPRINT Phase I [N = 24]) and in other identified clinical trials in adult patients (N = 347)<sup>††</sup>**

MedDRA SOC	MedDRA Term	Overall Frequency (All CTCAE grades) NF1 paediatric pool <sup>‡</sup> (N = 74)	Frequency of CTCAE grade 3 and Above <sup>†</sup> NF1 paediatric pool <sup>‡</sup> (N = 74)
Eye disorders	Vision blurred <sup>^</sup>	Very common (15%)	-
	Retinal pigment epithelial detachment (RPED)/ Central serous retinopathy (CSR)* <sup>††</sup>	Uncommon (0.6%)	-
	Retinal vein occlusion (RVO)* <sup>††</sup>	Uncommon (0.3%)	-

MedDRA SOC	MedDRA Term	Overall Frequency (All CTCAE grades)  NF1 paediatric pool <sup>‡</sup> (N = 74)	Frequency of CTCAE grade 3 and Above <sup>†</sup>  NF1 paediatric pool <sup>‡</sup> (N = 74)
Respiratory, thoracic & mediastinal disorders	Dyspnoea*	Common (8%)	-
Gastrointestinal disorders	Vomiting <sup>^</sup>	Very common (86%)	Common (9%)
	Diarrhoea <sup>^</sup>	Very common (81%)	Very common (15%)
	Nausea <sup>^</sup>	Very common (77%)	Common (3%)
	Stomatitis <sup>^</sup>	Very common (55%)	Common (1%)
	Dry mouth	Common (5%)	-
Skin and subcutaneous tissue disorders	Dry skin	Very common (65%)	Common (1%)
	Dermatitis acneiform <sup>^</sup>	Very common (61%)	Common (4%)
	Paronychia <sup>^</sup>	Very common (57%)	Very common (14%)
	Rashes (non- acneiform) <sup>^ *</sup>	Very common (53%)	Common (3%)
	Hair changes <sup>^ *</sup>	Very common (39%)	-
General disorders	Pyrexia	Very common (61%)	Common (8%)
	Asthenic events*	Very common (59%)	-
	Peripheral oedema*	Very common (31%)	-
	Facial oedema*	Common (8%)	-
Investigations	Blood CPK increased <sup>^</sup>	Very common (77%)	Common (9%)
	Haemoglobin decreased*	Very common (54%)	Common (3%)
	Hypoalbuminaemia	Very common (51%)	-
	AST increased	Very common (51%)	Common (1%)
	ALT increased	Very common (39%)	Common (3%)
	Blood creatinine increased	Very common (32%)	Common (1%)
	Ejection fraction decreased <sup>^</sup>	Very common (28%)	Common (1%)
	Increased blood pressure*	Very common (18%)	-

Per National Cancer Institute CTCAE version 4.03

CPK = creatine phosphokinase; AST = aspartate aminotransferase; ALT = alanine aminotransferase

<sup>^</sup> See Description of selected adverse reactions

<sup>†</sup> All reactions were CTCAE grade 3, except for one CTCAE grade 4 event of blood CPK increased and one CTCAE grade 4 event of blood creatinine increased. There were no deaths.

<sup>††</sup> Identified ADRs from other clinical trial experience in adult patients (N = 347), with multiple tumour types, receiving treatment with selumetinib (75 mg twice daily). These ADRs have not been reported in paediatric population with NF1 who have inoperable PN.

<sup>‡</sup> Paediatric pool (N = 74) percentage rounded to the nearest decimal.

\*ADRs based on grouping of individual preferred terms (PT):

Asthenic events: asthenia, fatigue

CSR/RPED: Detachment of macular retinal pigment epithelium, chorioretinopathy

Dyspnoea: dyspnoea exertional, dyspnoea, dyspnoea at rest

Facial oedema: face odema, periorbital oedema

Haemoglobin decreased: anaemia, haemoglobin decreased  
Hair changes: alopecia, hair colour change  
Increased blood pressure: blood pressure increased, hypertension  
Peripheral oedema: oedema peripheral, oedema, localised oedema, peripheral swelling  
Rashes (non-acneiform): rash pruritic, rash maculo-papular, rash papular, rash, rash erythematous, rash macular  
RVO: retinal vascular disorder, retinal vein occlusion, retinal vein thrombosis

### Description of selected adverse reactions

#### Left ventricular ejection fraction (LVEF) reduction

In SPRINT, Phase II Stratum 1, LVEF reduction (PT: ejection fraction decreased) was reported in 13 (26%) patients; all cases were grade 2, asymptomatic and did not lead to discontinuation; one (2%) case led to dose interruption then reduction. Of the 13 patients, 11 patients recovered and for 2 patients the outcome was not reported. The median time to first occurrence of LVEF reduction was 232 days (median duration 252 days). The majority of LVEF reduction adverse reactions were reported as reductions from baseline ( $\geq 10\%$  reduction) but were considered to remain in the normal range. Patients with LVEF lower than the institutional LLN at baseline were not included in the pivotal study. In addition, a small number of serious cases of LVEF reduction associated with selumetinib have been reported in paediatric patients who participated in an expanded access program. For clinical management of LVEF reduction (see sections 4.2 and 4.4).

#### Ocular toxicity

In SPRINT, Phase II Stratum 1, grade 1 and 2 adverse reactions of blurred vision were reported in 7 (14%) patients. Two patients required dose interruption. All adverse reactions were managed without dose reduction. For clinical management of new visual disturbances (see sections 4.2 and 4.4).

In addition, a single event of RPED was reported in a paediatric patient receiving selumetinib monotherapy (25 mg/m<sup>2</sup> twice daily) for pilocytic astrocytoma involving the optic pathway in an externally sponsored paediatric study (see sections 4.2 and 4.4).

#### Paronychia

In SPRINT, Phase II Stratum 1, paronychia was reported in 28 (56%) patients, the median time to first onset of maximum grade paronychia adverse reaction was 423 days and the median duration of adverse reactions was 51 days. The majority of these adverse reactions were grade 1 or 2 and were treated with supportive or symptomatic therapy and/or dose modification. Grade  $\geq 3$  events occurred in 4 (8%) patients. Ten patients (3 with a maximum grade 3 adverse reaction and 7 with a maximum grade 2 adverse reaction) had a selumetinib dose interruption for adverse reactions of paronychia, of whom 5 had dose interruption followed by dose reduction (2 patients required a second dose reduction). In one patient (2%) the event led to discontinuation.

#### Blood creatine phosphokinase (CPK) increase

Adverse reactions of blood CPK elevation occurred in 39 (78%) of patients in SPRINT Phase II Stratum 1. The median time to first onset of the maximum grade CPK increase was 112 days and the median duration of adverse reactions was 153 days. The majority of adverse reactions were grade 1 or 2 and resolved with no change in selumetinib dose. Grade  $\geq 3$  adverse reactions occurred in 3 (6%) patients. A grade 4 adverse reaction led to treatment interruption followed by dose reduction.

#### Gastrointestinal toxicities

In SPRINT, Phase II Stratum 1, vomiting (43 patients, 86%, median duration 3 days), diarrhoea (37 patients, 74%, median duration 6 days), nausea (36 patients, 72%, median duration 15 days), and stomatitis (26 patients, 52%, median duration 27 days) were the most commonly reported gastrointestinal (GI) reactions. The majority of these cases were grade 1 or 2 and did not require any dose interruptions or dose reductions.

Grade 3 adverse reactions were reported for diarrhoea (8 patients, 16%), nausea (2 patients, 4%), and vomiting (4 patients, 8%). For one patient diarrhoea led to dose reduction and subsequent

discontinuation. No dose reduction or discontinuation was required for adverse reactions of nausea, vomiting or stomatitis.

#### Skin toxicities

In SPRINT, Phase II Stratum 1, dermatitis acneiform was observed in 28 (56%) patients (median time to onset 43 days; median duration of 202 days for the maximum CTCAE grade event). The majority of these cases were grade 1 or 2, observed in post-pubertal patients (> 12 years) and did not require any dose interruptions or reductions. Grade 3 adverse reactions were reported in 3 (6%) patients.

Other (non-acneiform) rashes were observed in 27 (54%) patients in the pivotal study and were predominantly grade 1 or 2.

#### Hair changes

In SPRINT, Phase II Stratum 1, 16 (32%) of patients experienced hair changes (reported as hair lightening [PT: hair colour changes] in 12 patients (24%) and hair thinning [PT: alopecia] in 12 patients (24%)); in 8 patients (16%) both alopecia and hair colour changes were reported during treatment. All cases were grade 1 and did not require dose interruption or dose reduction.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

### **4.9 Overdose**

There is no specific treatment for overdose. If overdose occurs, patients should be closely monitored for signs and symptoms of adverse reactions and treated supportively with appropriate monitoring as necessary. Dialysis is ineffective in the treatment of overdose.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antineoplastic agents, protein kinase inhibitor, ATC code: L01EE04

#### Mechanism of action

Selumetinib is a selective inhibitor of mitogen activated protein kinase kinases 1 and 2 (MEK 1/2). Selumetinib blocks MEK activity and the RAF-MEK-ERK pathway. Therefore, MEK inhibition can block the proliferation and survival of tumour cells in which the RAF-MEK-ERK pathway is activated.

#### Clinical efficacy

The efficacy of Koselugo was evaluated in an open-label, multi-centre, single-arm study (SPRINT) Phase II Stratum 1 of 50 paediatric patients with NF1 inoperable PN that caused significant morbidity. Inoperable PN was defined as a PN that could not be surgically completely removed without risk for substantial morbidity due to encasement of, or close proximity to, vital structures, invasiveness, or high vascularity of the PN. Patients were excluded for the following ocular toxicities: any current or past history of CSR, current or past history of RVO, known intraocular pressure > 21 mmHg (or upper limit of normal adjusted by age) or uncontrolled glaucoma. Patients received 25 mg/m<sup>2</sup> (BSA) twice daily, for 28 days (1 treatment cycle), on a continuous dosing schedule. Treatment was discontinued if a patient was no longer deriving clinical benefit, experienced unacceptable toxicity or PN progression, or at the discretion of the investigator.

The target PN, the PN that caused relevant clinical symptoms or complications (PN-related morbidities), was evaluated for response rate using centrally read volumetric magnetic resonance

imaging (MRI) analysis per Response Evaluation in Neurofibromatosis and Schwannomatosis (REiNS) criteria. Tumour response was evaluated at baseline and while on treatment after every 4 cycles for 2 years, and then every 6 cycles.

Patients had target PN MRI volumetric evaluations and clinical outcome assessments, which included functional assessments and patient reported outcomes.

At enrolment, the median age of the patients was 10.2 years (range: 3.5 to 17.4 years), 60% were male and 84% were Caucasian.

The median target PN volume at baseline was 487.5 mL (range: 5.6 - 3820 mL). PN-related morbidities that were present in  $\geq 20\%$  of patients included disfigurement, motor dysfunction, pain, airway dysfunction, visual impairment, and bladder/bowel dysfunction.

The primary efficacy endpoint was objective response rate (ORR), defined as the percentage of patients with complete response (defined as disappearance of the target PN) or confirmed partial response (defined as  $\geq 20\%$  reduction in PN volume, confirmed at a subsequent tumour assessment within 3-6 months), based on National Cancer Institute (NCI) centralised review. Duration of response (DoR) was also evaluated.

Efficacy results are provided based on a data cut-off of March 2021, unless stated otherwise.

**Table 6. Efficacy results from SPRINT Phase II Stratum 1**

Efficacy parameter	SPRINT (N = 50)
<b>Objective response rate<sup>a, b</sup></b>	
Objective response rate, % (95% CI)	34 (68%) (53.3 - 80.5)
Complete response	0
Confirmed partial response, n (%) <sup>b</sup>	34(68%)
<b>Duration of response</b>	
DoR $\geq 12$ months, n (%)	31 (91.2%)
DoR $\geq 24$ months, n (%)	26 (76.5%)
DoR $\geq 36$ months, n (%)	21 (61.8%)

CI – confidence interval, DoR – duration of response.

<sup>a</sup> Responses required confirmation at least 3 months after the criteria for first partial response were met.

<sup>b</sup> Complete response: disappearance of the target lesion; partial response: decrease in target PN volume by  $\geq 20\%$  compared to baseline.

An independent centralized review of tumour response per REiNS criteria (data cut-off June 2018) resulted in an ORR of 44% (95% CI: 30.0, 58.7).

The median time to onset of response was 7.2 months (range: 3.3 months to 3.2 years). The median (min-max) time to the maximal PN shrinkage from baseline was 15.1 months (range: 3.3 months to 5.2 years). The median DoR from onset of response was not reached; at the time of data cut-off the median follow-up time was 41.3 months. The median time from treatment initiation to disease progression while on treatment was not reached.

At the time of data cut-off or last scan on treatment for patients who had discontinued treatment, 25 (50%) patients remained in confirmed partial response, 1 (2%) had unconfirmed partial responses, 12 (24%) had stable disease and 10 (20%) had progressive disease.

#### Paediatric population

The European Medicines Agency has deferred the obligation to submit the results of studies with Koselugo in one or more subsets of the paediatric population in NF1 PN (see section 4.2 for information on paediatric use).

This medicinal product has been authorised under a so-called “conditional approval” scheme. This means that further evidence on this medicinal product is awaited. The European Medicines Agency will review new information on this medicinal product at least every year and this SmPC will be updated as necessary.

## 5.2 Pharmacokinetic properties

At the recommended dose of 25 mg/m<sup>2</sup> twice daily in paediatric patients (3 to ≤ 18 years old), the geometric mean (coefficient of variation [CV%]) maximum plasma concentration (C<sub>max</sub>) was 731 (62%) ng/mL and that of the area under the plasma drug concentration curve (AUC<sub>0-12</sub>) following the first dose was 2009 (35%) ng·h/mL. Minimal accumulation of ~1.1-fold was observed at steady state upon twice daily dosing.

In paediatric patients, at a dose level of 25 mg/m<sup>2</sup>, selumetinib has an apparent oral clearance of 8.8 L/h, mean apparent volume of distribution at steady state of 78 L and mean elimination half-life of ~6.2 hours.

### Absorption

In healthy adult subjects, the mean absolute oral bioavailability of selumetinib was 62%. Following oral dosing, selumetinib is rapidly absorbed, producing peak steady state plasma concentrations (T<sub>max</sub>) between 1-1.5 hours post-dose.

### Effect of food

In separate clinical studies, in healthy adult subjects and in adult patients with advanced solid malignancies at a dose of 75 mg, co-administration of selumetinib with a high-fat meal resulted in a mean decrease in C<sub>max</sub> of 50% and 62%, respectively, compared to fasting administration. Selumetinib mean AUC was reduced by 16% and 19%, respectively, and the time to reach maximum concentration (T<sub>max</sub>) was delayed by approximately 1.5 to 3 hours (see section 4.2).

In healthy adult subjects at a dose of 50 mg, co-administration of selumetinib with a low-fat meal resulted in 60% lower C<sub>max</sub> when compared to fasting administration. Selumetinib AUC was reduced by 38%, and the time to reach maximum concentration (T<sub>max</sub>) was delayed by approximately 0.9 hours (see section 4.2).

### Distribution

The mean apparent volume of distribution at steady state of selumetinib across 20 to 30 mg/m<sup>2</sup> ranged from 78 to 171 L in paediatric patients, indicating moderate distribution into tissue.

*In vitro* plasma protein binding is 98.4% in humans. Selumetinib mostly binds to serum albumin (96.1%) than α-1 acid glycoprotein (< 35%).

### Biotransformation

*In vitro*, selumetinib undergoes phase 1 metabolic reactions including oxidation of the side chain, N-demethylation, and loss of the side chain to form amide and acid metabolites. CYP3A4 is the predominant isoform responsible for selumetinib oxidative metabolism with CYP2C19, CYP2C9, CYP2E1 and CYP3A5 involved to a lesser extent. *In vitro* studies indicate that selumetinib also undergoes direct phase 2 metabolic reactions to form glucuronide conjugates principally involving the enzymes UGT1A1 and UGT1A3. Glucuronidation is a significant route of elimination for selumetinib phase 1 metabolites involving several UGT isoforms.

Following oral dosing of <sup>14</sup>C-selumetinib to healthy male subjects, unchanged selumetinib (~40% of the radioactivity) with other metabolites including glucuronide of imidazoindazole metabolite (M2; 22%), selumetinib glucuronide (M4; 7%), N-desmethyl selumetinib (M8; 3%), and N-desmethyl carboxylic acid (M11; 4%) accounted for the majority of the circulating radioactivity in human plasma. N-desmethyl selumetinib represents less than 10% of selumetinib levels in human plasma but is approximately 3 to 5 times more potent than the parent compound, contributing to about 21% to 35% of the overall pharmacologic activity.

### Interactions

*In vitro*, selumetinib is not an inhibitor of CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, CYP3A4 and CYP2E1. *In vitro*, selumetinib is not an inducer of CYP1A2 and CYP2B6. Selumetinib is an inducer of CYP3A4 *in vitro*, this is however not expected to be clinically relevant.

*In vitro*, selumetinib inhibits UGT1A3, UGT1A4, UGT1A6 and UGT1A9 however these effects are not expected to be clinically relevant.

### Interactions with transport proteins

Based on *in vitro* studies, selumetinib is a substrate for BCRP and P-gp transporters but is unlikely to be subjected to clinically relevant drug interactions. *In vitro* studies suggest that selumetinib does not inhibit the breast cancer resistance protein (BCRP), P-glycoprotein (P-gp), OATP1B1, OATP1B3, OCT2, OAT1, MATE1 and MATE2K at the recommended paediatric dose. A clinically relevant effect on the pharmacokinetics of concomitantly administered substrates of OAT3 cannot be excluded.

### Elimination

In healthy adult subjects, following a single oral 75 mg dose of radiolabelled selumetinib, 59% of the dose was recovered in faeces (19% unchanged) while 33% of the administered dose (< 1% as parent) was found in urine by 9 days of sample collection.

### Special populations

#### Renal impairment

The exposure of 50 mg oral selumetinib was investigated in adult subjects with normal renal function (n = 11) and subjects with ESRD (n = 12). The ESRD group showed 16% and 28% lower C<sub>max</sub> and AUC, respectively, with the fraction of unbound selumetinib being 35% higher in ESRD subjects. As a result, the unbound C<sub>max</sub> and AUC ratios were 0.97 and 1.13 in the ESRD group when compared to the group with normal renal function. A small increase, approximately 20% AUC, in the N-desmethyl metabolite to parent ratio was detected in the ESRD group when compared to the normal group. As exposure in ESRD subjects was similar to those with normal renal function, investigations in mild, moderate and severe renally impaired subjects were not performed. Renal impairment is expected to have no meaningful influence on the exposure of selumetinib (see section 4.2).

#### Hepatic impairment

Adult subjects with normal hepatic function (n = 8) and mild hepatic impairment (Child-Pugh A, n = 8) were dosed with 50 mg selumetinib, subjects with moderate hepatic impairment (Child-Pugh B, n = 8) were administered a 50 or 25 mg dose, and subjects with severe hepatic impairment (Child-Pugh C, n = 8) were administered a 20 mg dose. Selumetinib total dose normalised AUC and unbound AUC were 86% and 69% respectively, in mild hepatic impairment patients, compared to the AUC values for subjects with normal hepatic function. Selumetinib exposure (AUC) was higher in patients with moderate (Child-Pugh B) and severe (Child-Pugh C) hepatic impairment; the total AUC and unbound AUC values were 159% and 141% (Child-Pugh B) and 157% and 317% (Child-Pugh C), respectively, of subjects with normal hepatic function (see section 4.2). There was a trend of lower protein binding in subjects with severe hepatic impairment although the protein binding remained > 99% (see section 4.3).

#### Ethnicity

Following a single-dose, selumetinib exposure appears to be higher in Japanese, non-Japanese-Asian and Indian healthy adult subjects compared to Western adult subjects, however, there is considerable overlap with Western subjects when corrected for body weight or BSA (see section 4.2).

#### Adult patients (> 18 years old)

The PK parameters in adult healthy subjects and adult patients with advanced solid malignancies, are similar to those in paediatric patients (3 to ≤ 18 years old) with NF1.

In adult patients,  $C_{max}$  and AUC increased dose proportionally over a 25 mg to 100 mg dose range.

### 5.3 Preclinical safety data

#### Genotoxicity

Selumetinib was positive in the mouse micronucleus study via an aneugenic mode of action. The free mean exposure ( $C_{max}$ ) at the no observed effect level (NOEL) was approximately 27-times greater than clinical free exposure at the maximum recommended human dose (MRHD) of 25 mg/m<sup>2</sup>.

#### Carcinogenicity

Selumetinib was not carcinogenic in rats or transgenic mice.

#### Repeat-dose toxicity

In repeat-dose toxicity studies in mice, rats and monkeys, the main effects seen after selumetinib exposure were in the skin, GI tract and bones. Scabs associated with microscopic erosions and ulceration at a free exposure similar to the clinical exposure (free AUC) at the MRHD were seen in rats. Inflammatory and ulcerative GI tract findings associated with secondary changes in the liver and lymphoreticular system at free exposures approximately 28 times the clinical free exposure at the MRHD were observed in mice. Growth plate (physeal) dysplasia was seen in male rats dosed for up to 3 months with selumetinib at a free exposure 11 times the clinical free exposure at the MRHD. GI findings showed evidence of reversibility following a recovery period. Reversibility for skin toxicities and physeal dysplasia was not evaluated. Vascular engorgement of the corpus cavernosum of the bulbocavernosus muscle were observed in male mice in a 26-week study at a dose of 40 mg/kg/day (28 times the free AUC in humans at the MRHD) leading to significant urinary tract obstruction as well as inflammation and luminal hemorrhage of the urethra leading to early death in male mice.

#### Reproductive toxicology

Developmental and reproduction toxicity studies were conducted in mice. Fertility was not affected in male mice at up to 40 mg/kg/day (corresponding to 22-fold the free AUC in humans at the MRHD). In females, mating performance and fertility were not affected at up to 75 mg/kg/day, but a reversible decrease in the number of live foetuses was observed at this dose level; the NOAEL for effects on reproductive performance was 5 mg/kg/day (approximately 3.5-fold the free AUC in humans at the MRHD). A treatment-related increase in the incidence of external malformations (open eye, cleft palate) was reported in absence of maternal toxicity in embryofoetal development studies at > 5 mg/kg/day, and in the pre- and post-natal development study at  $\geq 1$  mg/kg/day (corresponding to 0.4-fold the free  $C_{max}$  in humans at the MRHD). The other treatment related effects observed at non-maternotoxic dose levels in these studies consisted of embryo-lethality and decreased foetal weight at  $\geq 25$  mg/kg/day (corresponding to 22-fold the free AUC in humans at the MRHD), reductions in post-natal pup growth and at weaning a lower number of pups met the pupil constriction criterion at 15 mg/kg/day (corresponding to 3.6-fold the free  $C_{max}$  in humans at the MRHD). Selumetinib and its active metabolite were excreted in the milk of lactating mice at concentrations approximately the same as those in plasma.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

#### Capsule content

Tocofersolan (Vitamin E polyethylene glycol succinate /D  $\alpha$ -tocopheryl polyethylene glycol succinate).

#### Capsule shell

##### *Koselugo 10 mg hard capsules*

Hypromellose (E464)

Carrageenan (E407)

Potassium chloride (E508)



Titanium dioxide (E171)  
Carnauba wax (E903)

Koselugo 25 mg hard capsules

Hypromellose (E464)  
Carrageenan (E407)  
Potassium chloride (E508)  
Titanium dioxide (E171)  
Indigo carmine aluminium lake (E132)  
Iron oxide yellow (E172)  
Carnauba wax (E903)  
Maize starch

Printing ink

Koselugo 10 mg hard capsules

Shellac glaze, standard (E904)  
Iron oxide black (E172)  
Propylene glycol (E1520)  
Ammonium hydroxide (E527)

Koselugo 25 mg hard capsules

Iron oxide red (E172)  
Iron oxide yellow (E172)  
Indigo carmine aluminium lake (E132)  
Carnauba wax (E903)  
Shellac, standard (E904)  
Glyceryl mono-oleate

## 6.2 Incompatibilities

Not applicable.

## 6.3 Shelf life

3 years.

## 6.4 Special precautions for storage

Do not store above 30 °C.  
Store in the original bottle in order to protect from moisture and light.  
Keep the bottle tightly closed.

## 6.5 Nature and contents of container

Koselugo 10 mg hard capsules

High-density polyethylene (HDPE) plastic bottle with white child-resistant polypropylene closure.

Koselugo 25 mg hard capsules

High-density polyethylene (HDPE) plastic bottle with blue child-resistant polypropylene closure.

Each bottle contains 60 hard capsules and a silica gel desiccant. Each carton contains one bottle.

## 6.6 Special precautions for disposal and other handling

Patients should be instructed not to remove the desiccant from the bottle.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/21/1552/001 10 mg hard capsules  
EU/1/21/1552/002 25 mg hard capsules

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 17 June 2021  
Date of latest renewal: 25 April 2022

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>

## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**
- E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE CONDITIONAL MARKETING AUTHORISATION**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer responsible for batch release

AstraZeneca AB  
Global External Sourcing (GES)  
Astraallén  
Gärtunaporten (B674:5)  
151 85, Södertälje  
Sweden

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2)

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

### **• Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in Article 9 of Regulation (EC) No 507/2006 and, accordingly, the marketing authorisation holder (MAH) shall submit PSURs every 6 months.

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

The marketing authorisation holder (MAH) shall submit the first PSUR for this product within 6 months following authorisation.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

### **• Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

## **E. SPECIFIC OBLIGATION TO COMPLETE POST-AUTHORISATION MEASURES FOR THE CONDITIONAL MARKETING AUTHORISATION**

This being a conditional marketing authorisation and pursuant to Article 14-a of Regulation (EC) No 726/2004, the MAH shall complete, within the stated timeframe, the following measures:

<b>Description</b>	<b>Due date</b>
<p>Non-interventional post-authorisation safety study (PASS): in order to confirm the long-term safety of selumetinib in the treatment of symptomatic, inoperable PN in paediatric patients with NF1 aged 3 years and above, the applicant will conduct and submit the results of a non-interventional PASS in patients with NF1 who have been prescribed at least one dose of selumetinib and who are aged 3 to <math>\leq 18</math> years at the start of selumetinib treatment. A nested cohort of patients aged <math>\geq 8</math> years old (and prior to attainment of Tanner Stage V [sexual maturity rating]) will be followed prospectively.</p> <p>The clinical study report will be submitted by:</p>	<p>31/03/2028</p>

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Koselugo 10 mg hard capsules  
selumetinib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each hard capsule contains 10 mg of selumetinib (as hydrogen sulfate).

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

60 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not remove desiccant.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C.  
Store in the original bottle in order to protect from moisture and light.



**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/21/1552/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

koselugo 10 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Koselugo 10 mg hard capsules  
selumetinib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each hard capsule contains 10 mg of selumetinib (as hydrogen sulfate).

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

60 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not remove desiccant.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C.  
Store in the original bottle in order to protect from moisture and light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/21/1552/001

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

Koselugo 25 mg hard capsules  
selumetinib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each hard capsule contains 25 mg of selumetinib (as hydrogen sulfate).

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

60 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not remove desiccant.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C.  
Store in the original bottle in order to protect from moisture and light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/21/1552/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

koselugo 25 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC  
SN  
NN

**PARTICULARS TO APPEAR ON THE IMMEDIATE PACKAGING**

**BOTTLE LABEL**

**1. NAME OF THE MEDICINAL PRODUCT**

Koselugo 25 mg hard capsules  
selumetinib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each hard capsule contains 25 mg of selumetinib (as hydrogen sulfate).

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Hard capsule

60 hard capsules

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

Do not remove desiccant.

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Do not store above 30 °C.  
Store in the original bottle in order to protect from moisture and light.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/21/1552/002

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

**17. UNIQUE IDENTIFIER – 2D BARCODE**

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

**B. PACKAGE LEAFLET**



## Package leaflet: Information for the patient

**Koselugo 10 mg hard capsules**

**Koselugo 25 mg hard capsules**

selumetinib

▼ This medicine is subject to additional monitoring. This will allow quick identification of new safety information. You can help by reporting any side effects you may get. See the end of section 4 for how to report side effects.

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What Koselugo is and what it is used for
2. What you need to know before you take Koselugo
3. How to take Koselugo
4. Possible side effects
5. How to store Koselugo
6. Contents of the pack and other information

### 1. What Koselugo is and what it is used for

#### What Koselugo is and how it works

Koselugo contains the active substance selumetinib.

Selumetinib is a type of medicine called a MEK inhibitor. It works by blocking certain proteins involved in the growth of tumour cells.

Koselugo is expected to shrink tumours that grow along nerves, called plexiform neurofibromas. These tumours are caused by a genetic condition called neurofibromatosis type 1 (NF1).

#### What Koselugo is used for

Koselugo is used to treat children aged 3 years and above with plexiform neurofibromas that cannot be completely removed by surgery.

If you have any questions about how Koselugo works or why this medicine has been prescribed for you, ask your doctor.

### 2. What you need to know before you take Koselugo

#### Do not take Koselugo:

- if you are allergic to selumetinib or any of the other ingredients of this medicine (listed in section 6)
- if you have severe liver disease

If you are not sure, talk to your doctor, pharmacist or nurse before taking Koselugo.

### **Warnings and precautions**

Talk to your doctor, pharmacist or nurse before and during your treatment with Koselugo:

- if you have eye problems
- if you have heart problems
- if you have liver problems
- if you take supplements containing vitamin E
- if you cannot swallow the capsule whole

If any of the above apply to you (or you are not sure) talk to your doctor, pharmacist or nurse before taking this medicine.

### **Eye problems**

Koselugo can cause eye problems (see section 4 ‘Possible side effects’). **Tell your doctor straight away** if you get blurred vision or any other changes to your sight during treatment. Your doctor should examine your eyes if you have any new or worsening problems with your sight while you are taking this medicine.

### **Heart problems**

Koselugo can lower the amount of blood pumped by your heart (see section 4 ‘Possible side-effects’). Your doctor will check how well your heart works before and during your treatment with Koselugo.

### **Liver problems**

Koselugo can increase the amount of some liver enzymes in your blood (see section 4 ‘Possible side effects’). Your doctor will do blood tests before and during treatment to check how well your liver is working.

### **Supplemental vitamin E**

Koselugo capsules contain vitamin E that may increase your risk of bleeding. This means you should tell your doctor if you take other medicines that increase your risk of bleeding such as:

- acetylsalicylic acid (also known as aspirin) for pain and inflammation
- anticoagulant medicines (blood thinners) such as warfarin or other medicines used for preventing blood clots
- supplements that may increase your risk of bleeding, such as vitamin E

### **Difficulty swallowing capsules**

Talk to your doctor if you think you might have difficulties swallowing the capsules whole (see section 3 ‘How to take Koselugo’).

### **Skin, nail and hair problems**

Koselugo can cause skin rash, nail infection or hair thinning or changes in hair colour (see section 4 ‘Possible side effects’). Tell your doctor if any of these symptoms trouble you during treatment.

### **Children under 3 years old**

Do not give Koselugo to children below 3 years of age. This is because it has not been studied in this age group.

### **Other medicines and Koselugo**

Tell your doctor or pharmacist if you are taking, have recently taken, or might take any other medicines. This includes herbal medicines, supplements and medicines obtained without a prescription.

Koselugo can affect the way some other medicines work. Also, some other medicines can affect the way Koselugo works. Tell your doctor if you are taking any of the following medicines:

- clarithromycin or erythromycin (used to treat bacterial infections)

- carbamazepine or phenytoin (used to treat seizures and epilepsy)
- digoxin (used to treat heart failure)
- fexofenadine (used to treat symptoms of allergy)
- fluconazole or itraconazole (used to treat fungal infections)
- ketoconazole (used to treat Cushing's syndrome)
- furosemide (used to treat fluid retention by increasing the amount of urine you pass)
- methotrexate (used to treat some types of cancer, psoriasis or rheumatoid arthritis)
- omeprazole (used to treat acid reflux or stomach ulcer)
- rifampicin (used to treat tuberculosis (TB) and some other bacterial infections)
- St. John's wort (*Hypericum perforatum*), a herbal medicine (used to treat mild depression and other conditions)
- ticlopidine (used to prevent blood clots)

Tell your doctor or pharmacist if you are taking or have recently taken any of the above or any other medicines, even those that are not prescribed.

### **Koselugo with food and drink**

Do not drink grapefruit juice while you are taking Koselugo because, it can affect the way the medicine works.

### **Pregnancy – information for women**

Koselugo is not recommended during pregnancy. It may cause harm to an unborn baby.

If you think you may be pregnant or are planning to have a baby, ask your doctor for advice before taking this medicine. Your doctor may ask you to take a pregnancy test before starting treatment.

You should not become pregnant while taking this medicine. If you are able to become pregnant, you must use effective contraception. See 'Contraception - information for women and men' below.

If you become pregnant during treatment, tell your doctor straight away.

### **Pregnancy – information for men**

If your partner becomes pregnant while you are taking this medicine, tell your doctor straight away.

### **Contraception – information for women and men**

If you are sexually active you should use effective contraception while you are taking this medicine and for at least 1 week after the last dose. It is not known whether Koselugo may interfere with how well hormonal contraceptives work. Please tell your doctor if you are taking a hormonal contraceptive, as your doctor may recommend the addition of a non-hormonal method of birth control.

### **Breast-feeding**

Do not breast-feed if you are taking Koselugo. It is not known if Koselugo passes into breast milk.

### **Driving and using machines**

Koselugo can cause side effects that affect your ability to drive or use machines. Do not drive or use machines if you feel tired or if you have problems with your vision (such as blurred vision).

## **3. How to take Koselugo**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

### **How much to take**

Your doctor will work out the correct dose for you based on your height and weight. The doctor will tell you how many capsules of Koselugo to take.

Your doctor may prescribe a lower dose if you have problems with your liver (hepatic impairment).

Your doctor may reduce your dose if you have certain side effects while you are taking Koselugo (see section 4 'Possible side effects') or the doctor may interrupt treatment or stop it permanently.

### **How to take**

- Take Koselugo twice a day, about 12 hours apart.
- Take the capsules on an empty stomach. This means that:
  - you must wait at least 2 hours before taking Koselugo after eating and
  - after taking Koselugo you must wait at least 1 hour before you eat.
- Swallow the capsules whole with water.
- Do not chew, dissolve, or open the capsules.
- If you have, or think you might have difficulty swallowing capsules whole, talk to your doctor before starting treatment.

### **If you are sick**

If you are sick (vomit) at any time after taking Koselugo, do not take an extra dose. Take the next dose at the normal time.

### **If you take more Koselugo than you should**

If you have taken more Koselugo than you should, contact your doctor or pharmacist immediately.

### **If you forget to take Koselugo**

What to do if you forget to take a dose of Koselugo depends on how long it is until your next dose.

- If it is more than 6 hours until your next dose, take the missed dose. Then take the next dose at the normal time.
- If it is less than 6 hours until your next dose, skip the missed dose. Then take the next dose at the normal time.

Do not take a double dose (two doses at the same time) to make up for a forgotten dose.

### **If you stop taking Koselugo**

Do not stop taking Koselugo unless your doctor tells you.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

## **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

### **Possible serious side effects**

#### Eye (vision) problems

Koselugo can cause eye problems. Tell your doctor straight away if you get blurred vision (a very common side effect that may affect more than 1 in 10 people) or any other changes to your sight during treatment. Your doctor may ask you to stop taking this medicine or send you to a specialist, if you develop symptoms that include:

- blurred vision
- loss of vision
- dark spots in your vision (floaters)
- other changes to your vision (such as reduced vision)

Tell your doctor straight away if you notice any of the serious side effects above.

## Other side effects

Tell your doctor or pharmacist if you notice any of the following side effects:

### Very common (may affect more than 1 in 10 people)

- being sick (vomiting), feeling sick (nausea)
- diarrhoea
- inflammation of the mouth (stomatitis)
- skin and nail problems - signs may include dry skin, rash, redness around the fingernails
- hair thinning (alopecia), hair colour change
- feeling tired, weak or lacking energy
- fever (pyrexia)
- swelling of the hands or feet (peripheral oedema)
- a slight decrease in the amount of blood that the heart is pumping (ejection fraction decreased) – signs may include shortness of breath or swelling in your legs, ankles or feet
- high blood pressure (hypertension)
- reduced level of albumin, an essential protein in the blood (shown in blood tests)
- reduced haemoglobin, the oxygen-carrying protein in red blood cells (shown in blood tests)
- increase in enzymes (shown in blood tests) suggesting stress on the liver, kidney injury or muscle breakdown

### Common (may affect up to 1 in 10 people)

- dry mouth
- swelling of the face (facial oedema)
- shortness of breath (dyspnoea)

## Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## 5. How to store Koselugo

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the bottle and carton after EXP. The expiry date refers to the last day of that month.

Do not store above 30 °C.

Store in the original bottle in order to protect from moisture and light.

Keep the bottle tightly closed.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## 6. Contents of the pack and other information

### What Koselugo contains

The active substance is selumetinib. Each Koselugo 10 mg hard capsule contains 10 mg of selumetinib (as hydrogen sulfate). Each Koselugo 25 mg hard capsule contains 25 mg of selumetinib (as hydrogen sulfate).

The other ingredients in Koselugo 10 mg hard capsules are:

- capsule fill: vitamin E polyethylene glycol succinate (D  $\alpha$ -tocopheryl polyethylene glycol succinate).
- capsule shell: hypromellose (E464), carrageenan (E407), potassium chloride (E508), titanium dioxide (E171), carnauba wax (E903).
- printing ink: shellac standard (E904), iron oxide black (E172), propylene glycol (E1520) ammonium hydroxide (E527).

The other ingredients in Koselugo 25 mg hard capsules are:

- capsule fill: vitamin E polyethylene glycol succinate (D  $\alpha$ -tocopheryl polyethylene glycol succinate).
- capsule shell: hypromellose (E464), carrageenan (E407), potassium chloride (E508), titanium dioxide (E171), indigo carmine aluminium lake (E132), iron oxide yellow (E172), carnauba wax (E903), maize starch.
- printing ink: iron oxide red (E172), iron oxide yellow (E172), indigo carmine aluminium lake (E132), carnauba wax (E903), shellac, standard (E904), glyceryl mono-oleate.

### **What Koselugo looks like and contents of the pack**

Koselugo 10 mg hard capsule is a white to off-white, opaque, hard capsule which has a centre band and is marked with “SEL 10” in black ink.

Koselugo 25 mg hard capsule is a blue, opaque, hard capsule which has a centre band and is marked with “SEL 25” in black ink.

Koselugo is provided in white plastic bottles, capped with a white (10 mg) or blue (25 mg) child-resistant closure containing 60 hard capsules and a silica gel desiccant. Do not remove the desiccant from the bottle and do not swallow it.

### **Marketing Authorisation Holder**

AstraZeneca AB  
SE-151 85 Södertälje  
Sweden

### **Manufacturer**

AstraZeneca AB  
Global External Sourcing (GES)  
Astraallén  
Gärtunaporten  
SE-151 85 Södertälje  
Sweden

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Alexion Pharma Belgium  
Tel: +32 800 200 31

#### **Lietuva**

UAB AstraZeneca Lietuva  
Tel: +370 5 2660550

#### **България**

АстраЗенека България ЕООД  
Тел.: +359 24455000

#### **Luxembourg/Luxemburg**

Alexion Pharma Belgium  
Tél/Tel: +32 800 200 31

#### **Česká republika**

AstraZeneca Czech Republic s.r.o.  
Tel: +420 222 807 111

#### **Magyarország**

AstraZeneca Kft.  
Tel.: +36 1 883 6500

**Danmark**

AstraZeneca A/S  
Tlf: +45 43 66 64 62

**Deutschland**

AstraZeneca GmbH  
Tel: +49 40 809034100

**Eesti**

AstraZeneca  
Tel: +372 6549 600

**Ελλάδα**

AstraZeneca A.E.  
Τηλ: +30 210 6871500

**España**

Alexion Pharma Spain, S.L.  
Tel: +34 93 272 30 05

**France**

Alexion Pharma France SAS  
Tél: +33 1 47 32 36 21

**Hrvatska**

AstraZeneca d.o.o.  
Tel: +385 1 4628 000

**Ireland**

AstraZeneca Pharmaceuticals (Ireland)  
DAC  
Tel: +353 1609 7100

**Ísland**

Vistor hf.  
Sími: +354 535 7000

**Italia**

Alexion Pharma Italy srl  
Tel: +39 02 7767 9211

**Κύπρος**

Αλέκτωρ Φαρμακευτική Λτδ  
Τηλ: +357 22490305

**Latvija**

SIA AstraZeneca Latvija  
Tel: +371 67377100

**Malta**

Associated Drug Co. Ltd  
Tel: +356 2277 8000

**Nederland**

AstraZeneca BV  
Tel: +31 79 363 2222

**Norge**

AstraZeneca AS  
Tlf: +47 21 00 64 00

**Österreich**

AstraZeneca Österreich GmbH  
Tel: +43 1 711 31 0

**Polska**

AstraZeneca Pharma Poland Sp. z o.o.  
Tel.: +48 22 245 73 00

**Portugal**

Alexion Pharma Spain, S.L. - Sucursal em  
Portugal  
Tel: +34 93 272 30 05

**România**

AstraZeneca Pharma SRL  
Tel: +40 21 317 60 41

**Slovenija**

AstraZeneca UK Limited  
Tel: +386 1 51 35 600

**Slovenská republika**

AstraZeneca AB, o.z.  
Tel: +421 2 5737 7777

**Suomi/Finland**

AstraZeneca Oy  
Puh/Tel: +358 10 23 010

**Sverige**

AstraZeneca AB  
Tel: +46 8 553 26 000

**United Kingdom (Northern Ireland)**

AstraZeneca UK Ltd  
Tel: +44 1582 836 836

**This leaflet was last revised in**

This medicine has been given ‘conditional approval’. This means that there is more evidence to come about this medicine. The European Medicines Agency will review new information on this medicine at least every year and this leaflet will be updated as necessary.

**Other sources of information**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<http://www.ema.europa.eu>