

ANNEX I

SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Comtan 200 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 200 mg entacapone.

Excipients with known effect

Each tablet contains 1.82 mg sucrose, and 7.3 mg sodium as a constituent of the excipients.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet

Brownish-orange, oval, biconvex film-coated tablet with “Comtan” engraved on one side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Entacapone is indicated as an adjunct to standard preparations of levodopa/benserazide or levodopa/carbidopa for use in adult patients with Parkinson’s disease and end-of-dose motor fluctuations, who cannot be stabilised on those combinations.

4.2 Posology and method of administration

Entacapone should only be used in combination with levodopa/benserazide or levodopa/carbidopa. The prescribing information for these levodopa preparations is applicable to their concomitant use with entacapone.

Posology

One 200 mg tablet is taken with each levodopa/dopa decarboxylase inhibitor dose. The maximum recommended dose is 200 mg ten times daily, i.e. 2,000 mg of entacapone.

Entacapone enhances the effects of levodopa. Hence, to reduce levodopa-related dopaminergic adverse reactions, e.g. dyskinesias, nausea, vomiting and hallucinations, it is often necessary to adjust levodopa dosage within the first days to first weeks after initiating entacapone treatment. The daily dose of levodopa should be reduced by about 10–30% by extending the dosing intervals and/or by reducing the amount of levodopa per dose, according to the clinical condition of the patient.

If entacapone treatment is discontinued, it is necessary to adjust the dosing of other antiparkinsonian treatments, especially levodopa, to achieve a sufficient level of control of the parkinsonian symptoms.

Entacapone increases the bioavailability of levodopa from standard levodopa/benserazide preparations slightly (5–10%) more than from standard levodopa/carbidopa preparations. Hence, patients who are taking standard levodopa/benserazide preparations may need a larger reduction of levodopa dose when entacapone is initiated.

Renal impairment

Renal insufficiency does not affect the pharmacokinetics of entacapone and there is no need for dose adjustment. However, for patients who are receiving dialysis therapy, a longer dosing interval may be considered (see section 5.2).

Hepatic impairment

See section 4.3.

Elderly (≥ 65 years)

No dosage adjustment of entacapone is required for older people.

Paediatric population

The safety and efficacy of Comtan in children below age 18 have not been established. No data are available.

Method of administration

Entacapone is administered orally and simultaneously with each levodopa/carbidopa or levodopa/benserazide dose.

Entacapone can be taken with or without food (see section 5.2).

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Hepatic impairment.
- Pheochromocytoma.
- Concomitant use of entacapone and non-selective monoamine oxidase (MAO-A and MAO-B) inhibitors (e.g. phenelzine, tranylcypromine).
- Concomitant use of a selective MAO-A inhibitor plus a selective MAO-B inhibitor and entacapone (see section 4.5).
- A previous history of neuroleptic malignant syndrome (NMS) and/or non-traumatic rhabdomyolysis.

4.4 Special warnings and precautions for use

Rhabdomyolysis secondary to severe dyskinesias or neuroleptic malignant syndrome (NMS) has been observed rarely in patients with Parkinson's disease.

NMS, including rhabdomyolysis and hyperthermia, is characterised by motor symptoms (rigidity, myoclonus, tremor), mental status changes (e.g. agitation, confusion, coma), hyperthermia, autonomic dysfunction (tachycardia, labile blood pressure) and elevated serum creatine phosphokinase. In individual cases, only some of these symptoms and/or findings may be evident.

Neither NMS nor rhabdomyolysis have been reported in association with entacapone treatment from controlled trials in which entacapone was discontinued abruptly. Since the introduction into the market, isolated cases of NMS have been reported, especially following abrupt reduction or discontinuation of entacapone and other concomitant dopaminergic medicinal products. When considered necessary, withdrawal of entacapone and other dopaminergic treatment should proceed slowly, and if signs and/or symptoms occur despite a slow withdrawal of entacapone, an increase in levodopa dosage may be necessary.

Entacapone therapy should be administered cautiously to patients with ischaemic heart disease.

Because of its mechanism of action, entacapone may interfere with the metabolism of medicinal

products containing a catechol group and potentiate their action. Thus, entacapone should be administered cautiously to patients being treated with medicinal products metabolised by catechol-O-methyl transferase (COMT), e.g. rimeterole, isoprenaline, adrenaline, noradrenaline, dopamine, dobutamine, alpha-methyldopa, and apomorphine (see also section 4.5).

Entacapone is always given as an adjunct to levodopa treatment. Hence, the precautions valid for levodopa treatment should also be taken into account for entacapone treatment. Entacapone increases the bioavailability of levodopa from standard levodopa/benserazide preparations 5–10% more than from standard levodopa/carbidopa preparations. Consequently, adverse dopaminergic reactions may be more frequent when entacapone is added to levodopa/benserazide treatment (see also section 4.8). To reduce levodopa-related dopaminergic adverse reactions, it is often necessary to adjust levodopa dosage within the first days to first weeks after initiating entacapone treatment, according to the clinical condition of the patient (see sections 4.2 and 4.8).

Entacapone may aggravate levodopa-induced orthostatic hypotension. Entacapone should be given cautiously to patients who are taking other medicinal products which may cause orthostatic hypotension.

In clinical studies, undesirable dopaminergic effects, e.g. dyskinesia, were more common in patients who received entacapone and dopamine agonists (such as bromocriptine), selegiline or amantadine compared to those who received placebo with this combination. The doses of other antiparkinsonian medicinal products may need to be adjusted when entacapone treatment is initiated.

Entacapone in association with levodopa has been associated with somnolence and episodes of sudden sleep onset in patients with Parkinson's disease and caution should therefore be exercised when driving or operating machines (see also section 4.7).

For patients experiencing diarrhoea, a follow-up of weight is recommended in order to avoid potential excessive weight decrease. Prolonged or persistent diarrhoea appearing during use of entacapone may be a sign of colitis. In the event of prolonged or persistent diarrhoea, the medicinal product should be discontinued and appropriate medical therapy and investigations considered.

Patients should be regularly monitored for the development of impulse control disorders. Patients and carers should be made aware that behavioural symptoms of impulse control disorders including pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists and/or other dopaminergic treatments such as Comtan in association with levodopa. Review of treatment is recommended if such symptoms develop.

For patients who experience progressive anorexia, asthenia and weight decrease within a relatively short period of time, a general medical evaluation including liver function should be considered.

Comtan tablets contain sucrose. Therefore, patients with rare hereditary problems of fructose intolerance, glucose-galactose malabsorption or sucrase-isomaltase insufficiency should not take this medicine.

This medicinal product contains 7.3 mg sodium per tablet. The maximum recommended daily dose (10 tablets) contains 73 mg sodium, equivalent to 4% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

4.5 Interaction with other medicinal products and other forms of interaction

No interaction of entacapone with carbidopa has been observed with the recommended treatment schedule. Pharmacokinetic interaction with benserazide has not been studied.

In single-dose studies in healthy volunteers, no interactions were observed between entacapone and

imipramine or between entacapone and moclobemide. Similarly, no interactions between entacapone and selegiline were observed in repeated-dose studies in parkinsonian patients. However, the experience of the clinical use of entacapone with several medicinal products, including MAO-A inhibitors, tricyclic antidepressants, noradrenaline reuptake inhibitors such as desipramine, maprotiline and venlafaxine, and medicinal products that are metabolised by COMT (e.g. catechol-structured compounds: rimiterol, isoprenaline, adrenaline, noradrenaline, dopamine, dobutamine, alpha-methyldopa, apomorphine, and paroxetine) is still limited. Caution should be exercised when these medicinal products are used concomitantly with entacapone (see also sections 4.3 and 4.4).

Entacapone may be used with selegiline (a selective MAO-B inhibitor), but the daily dose of selegiline should not exceed 10 mg.

Entacapone may form chelates with iron in the gastrointestinal tract. Entacapone and iron preparations should be taken at least 2–3 hours apart (see section 4.8).

Entacapone binds to human albumin binding site II which also binds several other medicinal products, including diazepam and ibuprofen. Clinical interaction studies with diazepam and non-steroidal anti-inflammatory medicinal products have not been carried out. According to *in vitro* studies, significant displacement is not anticipated at therapeutic concentrations of the medicinal products.

Due to its affinity to cytochrome P450 2C9 *in vitro* (see section 5.2), entacapone may potentially interfere with medicinal products with metabolism dependent on this isoenzyme, such as S-warfarin. However, in an interaction study in healthy volunteers, entacapone did not change the plasma levels of S-warfarin, while the AUC for R-warfarin increased on average by 18% [CI₉₀ 11–26%]. The INR values increased on average by 13% [CI₉₀ 6–19%]. Thus, control of INR is recommended when entacapone treatment is initiated for patients receiving warfarin.

4.6 Fertility, pregnancy and lactation

Pregnancy

No overt teratogenic or primary foetotoxic effects were observed in animal studies in which the exposure levels of entacapone were markedly higher than the therapeutic exposure levels. As there is no experience in pregnant women, entacapone should not be used during pregnancy.

Breast-feeding

In animal studies entacapone was excreted in milk. The safety of entacapone in infants is unknown. Women should not breast-feed during treatment with entacapone.

4.7 Effects on ability to drive and use machines

Comtan in association with levodopa may have a major influence on the ability to drive and use machines. Entacapone may, together with levodopa, cause dizziness and symptomatic orthostatism. Therefore, caution should be exercised when driving or using machines.

Patients being treated with entacapone in association with levodopa and presenting with somnolence and/or sudden sleep onset episodes must be instructed to refrain from driving or engaging in activities where impaired alertness may put themselves or others at risk of serious injury or death (e.g. operating machines) until such recurrent episodes have resolved (see also section 4.4).

4.8 Undesirable effects

Summary of the safety profile

The most frequent adverse reactions caused by entacapone relate to the increased dopaminergic activity and occur most commonly at the beginning of treatment. Reduction of levodopa dosage decreases the severity and frequency of these reactions. The other major class of adverse reactions are gastrointestinal symptoms, including nausea, vomiting, abdominal pain, constipation and diarrhoea. Urine may be discoloured reddish-brown by entacapone, but this is a harmless phenomenon.

Usually the adverse reactions caused by entacapone are mild to moderate. In clinical studies the most common adverse reactions leading to discontinuation of entacapone treatment have been gastrointestinal symptoms (e.g. diarrhoea, 2.5%) and increased dopaminergic adverse reactions of levodopa (e.g. dyskinesias, 1.7%).

Dyskinesias (27%), nausea (11%), diarrhoea (8%), abdominal pain (7%) and dry mouth (4.2%) were reported significantly more often with entacapone than with placebo in pooled data from clinical studies involving 406 patients taking the medicinal product and 296 patients taking placebo.

Some of the adverse reactions, such as dyskinesia, nausea, and abdominal pain, may be more common with the higher doses (1,400 to 2,000 mg per day) than with the lower doses of entacapone.

Tabulated list of adverse reactions

The following adverse reactions, listed below in Table 1, have been accumulated both from clinical studies with entacapone and since the introduction of entacapone into the market.

Table 1* Adverse drug reactions

Psychiatric disorders	Common:	Insomnia, hallucinations, confusion, paroniria
	Very rare:	Agitation
Nervous system disorders	Very common:	Dyskinesia
	Common:	Parkinsonism aggravated, dizziness, dystonia, hyperkinesia
Cardiac disorders**	Common:	Ischaemic heart disease events other than myocardial infarction (e.g. angina pectoris)
	Uncommon:	Myocardial infarction
Gastrointestinal disorders	Very common:	Nausea
	Common:	Diarrhoea, abdominal pain, dry mouth, constipation, vomiting
	Very rare:	Anorexia
	Not known:	Colitis
Hepatobiliary disorders	Rare:	Hepatic function tests abnormal
	Not known:	Hepatitis with mainly cholestatic features (see section 4.4)
Skin and subcutaneous tissue disorders	Rare:	Erythematous or maculopapular rash
	Very rare:	Urticaria
	Not known:	Skin, hair, beard and nail discolourations
Renal and urinary disorders	Very common:	Urine discolouration

General disorders and administration site conditions

Common:	Fatigue, sweating increased, fall
Very rare:	Weight decrease

- * Adverse reactions are ranked under headings of frequency, the most frequent first, using the following convention: Very common ($\geq 1/10$); common ($\geq 1/100$ to $< 1/10$); uncommon ($\geq 1/1,000$ to $< 1/100$); rare ($\geq 1/10,000$ to $< 1/1,000$); very rare ($< 1/10,000$), not known (cannot be estimated from the available data, since no valid estimate can be derived from clinical trials or epidemiological studies).
- ** The incidence rates of myocardial infarction and other ischaemic heart disease events (0.43% and 1.54%, respectively) are derived from an analysis of 13 double-blind studies involving 2,082 patients with end-of-dose motor fluctuations receiving entacapone.

Description of selected adverse reactions

Entacapone in association with levodopa has been associated with isolated cases of excessive daytime somnolence and sudden sleep onset episodes.

Impulse control disorders: Pathological gambling, increased libido, hypersexuality, compulsive spending or buying, binge eating and compulsive eating can occur in patients treated with dopamine agonists and/or other dopaminergic treatments such as Comtan in association with levodopa (see section 4.4).

Isolated cases of NMS have been reported following abrupt reduction or discontinuation of entacapone and other dopaminergic treatments.

Isolated cases of rhabdomyolysis have been reported.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

The post-marketing data include isolated cases of overdose in which the reported highest daily dose of entacapone has been 16,000 mg. The acute symptoms and signs in these cases of overdose included confusion, decreased activity, somnolence, hypotonia, skin discolouration and urticaria. Management of acute overdose is symptomatic.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: other dopaminergic agents, ATC code: N04BX02.

Entacapone belongs to a new therapeutic class, catechol-O-methyl transferase (COMT) inhibitors. It is a reversible, specific, and mainly peripherally acting COMT inhibitor designed for concomitant administration with levodopa preparations. Entacapone decreases the metabolic loss of levodopa to 3-O-methyldopa (3-OMD) by inhibiting the COMT enzyme. This leads to a higher levodopa AUC. The amount of levodopa available to the brain is increased. Entacapone thus prolongs the clinical response to levodopa.

Entacapone inhibits the COMT enzyme mainly in peripheral tissues. COMT inhibition in red blood cells closely follows the plasma concentrations of entacapone, thus clearly indicating the reversible nature of COMT inhibition.

Clinical studies

In two phase III double-blind studies in a total of 376 patients with Parkinson's disease and end-of-dose motor fluctuations, entacapone or placebo was given with each levodopa/dopa decarboxylase inhibitor dose. The results are given in Table 2. In study I, daily ON time (hours) was measured from home diaries and in study II, the proportion of daily ON time.

Table 2 Daily ON time (Mean ± SD)

Study I: Daily On time (h)			
	Entacapone (n=85)	Placebo (n=86)	Difference
Baseline	9.3±2.2	9.2±2.5	
Week 8-24	10.7±2.2	9.4±2.6	1 h 20 min (8.3%) CI _{95%} 45 min, 1 h 56
Study II: Proportion of daily On time (%)			
	Entacapone (n=103)	Placebo (n=102)	Difference
Baseline	60.0±15.2	60.8±14.0	
Week 8-24	66.8±14.5	62.8±16.80	4.5% (0 h 35 min) CI _{95%} 0.93%, 7.97%

There were corresponding decreases in OFF time.

The % change from baseline in OFF time was -24% in the entacapone group and 0% in the placebo group in study I. The corresponding figures in study II were -18% and -5%.

5.2 Pharmacokinetic properties

General characteristics of the active substance

Absorption

There are large intra- and interindividual variations in the absorption of entacapone.

The peak concentration (C_{max}) in plasma is usually reached about one hour after ingestion of a 200 mg entacapone tablet. The substance is subject to extensive first-pass metabolism. The bioavailability of entacapone is about 35% after an oral dose. Food does not affect the absorption of entacapone to any significant extent.

Distribution

After absorption from the gastrointestinal tract, entacapone is rapidly distributed to the peripheral tissues with a distribution volume of 20 litres at steady state (V_{dss}). Approximately 92% of the dose is eliminated during β -phase with a short elimination half-life of 30 minutes. The total clearance of entacapone is about 800 ml/min.

Entacapone is extensively bound to plasma proteins, mainly to albumin. In human plasma the unbound fraction is about 2.0% in the therapeutic concentration range. At therapeutic concentrations, entacapone does not displace other extensively bound substances (e.g. warfarin, salicylic acid, phenylbutazone, or diazepam), nor is it displaced to any significant extent by any of these substances at therapeutic or higher concentrations.

Biotransformation

A small amount of entacapone, the (*E*)-isomer, is converted to its (*Z*)-isomer. The (*E*)-isomer accounts for 95% of the AUC of entacapone. The (*Z*)-isomer and traces of other metabolites account for the remaining 5%.

Data from *in vitro* studies using human liver microsomal preparations indicate that entacapone inhibits cytochrome P450 2C9 (IC₅₀ ~4 µM). Entacapone showed little or no inhibition of other types of P450 isoenzymes (CYP1A2, CYP2A6, CYP2D6, CYP2E1, CYP3A and CYP2C19) (see section 4.5).

Elimination

The elimination of entacapone occurs mainly by non-renal metabolic routes. It is estimated that 80–90% of the dose is excreted in faeces, although this has not been confirmed in man. Approximately 10–20% is excreted in urine. Only traces of entacapone are found unchanged in urine. The major part (95%) of the product excreted in urine is conjugated with glucuronic acid. Of the metabolites found in urine only about 1% have been formed through oxidation.

Characteristics in patients

The pharmacokinetic properties of entacapone are similar in both young people and older people. The metabolism of the medicinal product is slowed in patients with mild to moderate liver insufficiency (Child-Pugh Class A and B), which leads to an increased plasma concentration of entacapone in both the absorption and elimination phases (see section 4.3). Renal impairment does not affect the pharmacokinetics of entacapone. However, a longer dosing interval may be considered for patients who are receiving dialysis therapy.

5.3 Preclinical safety data

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and carcinogenic potential. In repeated dose toxicity studies, anaemia most likely due to iron chelating properties of entacapone was observed. Regarding reproduction toxicity, decreased foetal weight and a slightly delayed bone development were noticed in rabbits at systemic exposure levels in the therapeutic range.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Microcrystalline cellulose
Mannitol
Croscarmellose sodium
Hydrogenated vegetable oil
Magnesium stearate

Film-coating

Hypromellose
Polysorbate 80
Glycerol 85%
Sucrose
Yellow iron oxide (E172)
Red iron oxide (E172)
Titanium dioxide (E171)

Magnesium stearate

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

6.5 Nature and contents of container

White high-density polyethylene (HDPE) bottles with white tamper proof polypropylene (PP) closures containing 30, 60 or 100 tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

No special requirements for disposal.

7. MARKETING AUTHORISATION HOLDER

Orion Corporation
Orionintie 1
FI-02200 Espoo
Finland

8. MARKETING AUTHORISATION NUMBER(S)

EU/1/98/081/005-007

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 22 September 1998

Date of latest renewal: 03 September 2008

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturer responsible for batch release

Orion Corporation Orion Pharma
Joensuunkatu 7
FI-24100 Salo
Finland

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to medical prescription.

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

Not applicable.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING

CARTON AND BOTTLE LABEL

1. NAME OF THE MEDICINAL PRODUCT

Comtan 200 mg film-coated tablets
entacapone

2. STATEMENT OF ACTIVE SUBSTANCE(S)

1 tablet contains 200 mg of entacapone.

3. LIST OF EXCIPIENTS

Contains sucrose and sodium.

4. PHARMACEUTICAL FORM AND CONTENTS

30 film-coated tablets
60 film-coated tablets
100 film-coated tablets

5. METHOD AND ROUTE(S) OF ADMINISTRATION

Read the package leaflet before use.
Oral use.

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING(S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

Carton

Orion Corporation
Orionintie 1
FI-02200 Espoo
Finland

Label

Orion Corporation

12. MARKETING AUTHORISATION NUMBER(S)

EU/1/98/081/005	30 film-coated tablets
EU/1/98/081/006	100 film-coated tablets
EU/1/98/081/007	60 film-coated tablets

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Comtan 200 mg [carton only]

17. UNIQUE IDENTIFIER – 2D BARCODE

[carton only]

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

[carton only]

PC {number}
SN {number}
<NN {number}>

B. PACKAGE LEAFLET

Package leaflet: Information for the user

Comtan 200 mg film-coated tablets entacapone

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Comtan is and what it is used for
2. What you need to know before you take Comtan
3. How to take Comtan
4. Possible side effects
5. How to store Comtan
6. Contents of the pack and other information

1. What Comtan is and what it is used for

Comtan tablets contain entacapone and are used together with levodopa to treat Parkinson's disease. Comtan aids levodopa in relieving the symptoms of Parkinson's disease. Comtan has no effect on relieving the symptoms of Parkinson's disease unless taken with levodopa.

2. What you need to know before you take Comtan

Do not take Comtan

- if you are allergic to entacapone or any of the other ingredients of this medicine (listed in section 6);
- if you have a tumour of the adrenal gland (known as pheochromocytoma; this may increase the risk of severe high blood pressure);
- if you are taking certain antidepressants (ask your doctor or pharmacist whether your antidepressive medicine can be taken together with Comtan);
- if you have liver disease;
- if you have ever suffered from a rare reaction to antipsychotic medicines called neuroleptic malignant syndrome (NMS). See Section 4 Possible side effects for the characteristics of NMS;
- if you have ever suffered from a rare muscle disorder called rhabdomyolysis which was not caused by injury.

Warnings and precautions

Talk to your doctor or pharmacist before taking Comtan:

- if you have ever had a heart attack or any other diseases of the heart;
- if you are taking a medicine which may cause dizziness or light-headedness (low blood pressure) when rising from a chair or bed;
- if you experience prolonged diarrhoea consult your doctor as it may be a sign of inflammation of the colon;
- if you experience diarrhoea, monitoring of your weight is recommended in order to avoid

- potentially excessive weight loss;
- if you experience increasing loss of appetite, weakness, exhaustion and weight loss within a relatively short period of time, a general medical evaluation including liver function should be considered.

Tell your doctor if you or your family/carer notices you are developing urges or cravings to behave in ways that are unusual for you or you cannot resist the impulse, drive or temptation to carry out certain activities that could harm yourself or others. These behaviours are called impulse control disorders and can include addictive gambling, excessive eating or spending, an abnormally high sex drive or a preoccupation with an increase in sexual thoughts or feelings. Your doctor may need to review your treatments.

As Comtan tablets will be taken together with other levodopa medicines, please also read the package leaflets of these medicines carefully.

The dose of other medicines to treat Parkinson's disease may need to be adjusted when you start taking Comtan. Follow the instructions that your doctor has given you.

Neuroleptic Malignant Syndrome (NMS) is a serious but rare reaction to certain medicines, and may occur especially when Comtan and other medicines to treat Parkinson's disease are suddenly stopped or the dose is suddenly reduced. For the characteristics of NMS see Section 4 Possible side effects. Your doctor may advise you to slowly discontinue the treatment with Comtan and other medicines to treat Parkinson's disease.

Comtan taken with levodopa may cause drowsiness and may cause you to sometimes suddenly fall asleep. If this happens, you should not drive or use any tools or machines (see "Driving and using machines").

Other medicines and Comtan

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. In particular please tell your doctor if you are taking any of the following:

- rimiterole, isoprenaline, adrenaline, noradrenaline, dopamine, dobutamine, alpha-methyldopa, apomorphine;
- antidepressants including desipramine, maprotiline, venlafaxine, paroxetine;
- warfarin used to thin the blood;
- iron supplements. Comtan may make it harder for you to digest iron. Therefore, do not take Comtan and iron supplements at the same time. After taking one of them, wait at least 2 to 3 hours before taking the other.

Pregnancy, breast-feeding and fertility

Do not use Comtan during pregnancy or if you are breast-feeding.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

Driving and using machines

Comtan taken together with levodopa may lower your blood pressure, which may make you feel light-headed or dizzy. Be particularly careful when you drive or when you use tools or machinery.

In addition, Comtan taken with levodopa may make you feel very drowsy, or cause you to sometimes suddenly fall asleep.

Do not drive or operate machinery if you experience these side effects.

Comtan contains sucrose and sodium

Comtan tablets contain a sugar called sucrose. If you have been told by your doctor that you have an

intolerance to some sugars, contact your doctor before taking this medicinal product.

This medicine contains 7.3 mg sodium (main component of cooking/table salt) in each tablet. The maximum recommended daily dose (10 tablets) contains 73 mg of sodium. This is equivalent to 4% of the recommended maximum daily dietary intake of sodium for an adult.

3. How to take Comtan


Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Comtan is taken together with medicines containing levodopa (either levodopa/carbidopa preparations or levodopa/benserazide preparations). You may also use other medicines to treat Parkinson's disease at the same time.

The recommended dose of Comtan is one 200 mg tablet with each levodopa dose. The maximum recommended dose is 10 tablets per day, i.e. 2,000 mg of Comtan.

If you are receiving dialysis for renal insufficiency, your doctor may tell you to increase the time between doses.

To open the bottle for the first time: open the closure, and then press with your thumb on the seal until it breaks. See picture 1.



Use in children and adolescents

Experience with Comtan in patients under 18 years is limited. Therefore, the use of Comtan in children or adolescents cannot be recommended.

If you take more Comtan than you should

In the event of an overdose, consult your doctor, pharmacist or the nearest hospital immediately.

If you forget to take Comtan

If you forget to take the Comtan tablet with your levodopa dose, you should continue the treatment by taking the next Comtan tablet with your next levodopa dose.

Do not take a double dose to make up for a forgotten tablet.

If you stop taking Comtan

Do not stop taking Comtan unless your doctor tells you to.

When stopping, your doctor may need to re-adjust the dosage of your other medicines to treat Parkinson's disease. Suddenly stopping Comtan and other medicines to treat Parkinson's disease may result in unwanted side effects. See Section 2 Warnings and precautions.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. Usually side effects caused by Comtan are mild to moderate.

Some of the side effects are often caused by the increased effects of levodopa therapy and are most common at the start of treatment. If you experience such effects at the start of treatment with Comtan you should contact your doctor who may decide to adjust your dosage of levodopa.

Very common (may affect more than 1 in 10 people):

- Uncontrollable movements with difficulty in performing voluntary movements (dyskinesias);
- feeling sick (nausea);
- harmless reddish-brown discolouration of urine.

Common (may affect up to 1 in 10 people):

- Excessive movements (hyperkinesias), worsening of symptoms of Parkinson's disease, prolonged muscle cramps (dystonia);
- being sick (vomiting), diarrhoea, abdominal pain, constipation, dry mouth;
- dizziness, tiredness, increased sweating, fall;
- hallucinations (seeing/hearing/feeling/smelling things that are not really there), sleeplessness, vivid dreams, and confusion;
- heart or artery disease events (e.g. chest pain).

Uncommon (may affect up to 1 in 100 people):

- Heart attack.

Rare (may affect up to 1 in 1,000 people):

- Rashes;
- abnormal results in liver function test.

Very rare (may affect up to 1 in 10,000 people):

- Agitation;
- decreased appetite, weight loss;
- hives.

Not known (frequency cannot be estimated from the available data):

- Inflammation of the colon (colitis), inflammation of the liver (hepatitis) with yellowing of the skin and whites of the eyes;
- discolouration of the skin, hair, beard and nails.

When Comtan is given at higher doses:

In doses of 1,400 to 2,000 mg per day, the following side effects are more common:

- Uncontrollable movements;
- nausea;
- abdominal pain.

Other important side effects which may occur:

- Comtan taken with levodopa may rarely make you feel very drowsy during the day, and cause you to suddenly fall asleep;
- Neuroleptic Malignant Syndrome (NMS) is a rare severe reaction to medicines used to treat disorders of the nervous system. It is characterised by stiffness, muscle twitching, shaking, agitation, confusion, coma, high body temperature, increased heart rate and unstable blood pressure;
- a rare severe muscle disorder (rhabdomyolysis) which causes pain, tenderness and weakness of the muscles and may lead to kidney problems.

You may experience the following side effects:

- Inability to resist the impulse to perform an action that could be harmful, which may include:
 - strong impulse to gamble excessively despite serious personal or family consequences.
 - altered or increased sexual interest and behaviour of significant concern to you or to others, for example, an increased sexual drive.
 - uncontrollable excessive shopping or spending.
 - binge eating (eating large amounts of food in a short time period) or compulsive eating (eating more food than normal and more than is needed to satisfy your hunger).

Tell your doctor if you experience any of these behaviours; they will discuss ways of managing or reducing the symptoms.

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Comtan

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the bottle label. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Comtan contains

- The active substance is entacapone. Each tablet contains 200 mg of entacapone.
- The other ingredients are microcrystalline cellulose, mannitol, croscarmellose sodium, hydrogenated vegetable oil, magnesium stearate.
- The film-coating contains hypromellose, polysorbate 80, glycerol 85%, sucrose, yellow iron oxide (E172), red iron oxide (E172) and titanium dioxide (E171), magnesium stearate.

What Comtan looks like and contents of the pack

Comtan 200 mg film-coated tablets are brownish-orange, oval tablets with “Comtan” engraved on one side. They are packed in bottles.

There are three different pack sizes (bottles containing 30, 60, or 100 tablets). Not all pack sizes may be marketed.

Marketing Authorisation Holder

Orion Corporation
Orionintie 1
FI-02200 Espoo
Finland

Manufacturer

Orion Corporation Orion Pharma
Joensuunkatu 7
FI-24100 Salo
Finland

For further information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

Orion Pharma BVBA/SPRL
Tél/Tel: +32 (0)15 64 10 20

България

Orion Corporation
Тел.: +358 10 4261

Česká republika

Orion Pharma s.r.o.
Tel: +420 234 703 305

Danmark

Orion Corporation
Tlf: +358 10 4261

Deutschland

Orion Corporation
Tel: +358 10 4261

Eesti

Orion Corporation
Tel: +358 10 4261

Ελλάδα

Orion Pharma Hellas M.E.Π.E
Τηλ: +30 210 980 3355

España

Orion Pharma S.L.
Tel: +34 91 599 86 01

France

Orion Pharma
Tél: +33 (0) 1 85 18 00 00

Hrvatska

Orion Pharma d.o.o.
Tel. +386 (0) 1 600 8015

Ireland

Orion Corporation
Tel: +358 10 4261

Lietuva

Orion Corporation
Tel: +358 10 4261

Luxembourg/Luxemburg

Orion Pharma BVBA/SPRL
Tél/Tel: +32 (0)15 64 10 20

Magyarország

Orion Pharma Kft.
Tel.: +36 1 239 9095

Malta

Orion Corporation
Tel: +358 10 4261

Nederland

Orion Pharma BVBA/SPRL
Tel: +32 (0)15 64 10 20

Norge

Orion Corporation
Tlf: +358 10 4261

Österreich

Orion Pharma GmbH
Tel: +49 40 899 6890

Polska

Orion Corporation
Tel.: +358 10 4261

Portugal

Orionfín Unipessoal Lda
Tel: +351 21 154 68 20

România

Orion Corporation
Tel: +358 10 4261

Slovenija

Orion Corporation
Tel: +358 10 4261

Ísland

Orion Corporation
Sími: +358 10 4261

Italia

Orion Pharma S.r.l.
Tel: +39 02 67876111

Κύπρος

Orion Corporation
Τηλ: +358 10 4261

Latvija

Orion Corporation
Tel: +358 10 4261

Slovenská republika

Orion Pharma s.r.o
Tel: +420 234 703 305

Suomi/Finland

Orion Corporation
Puh/Tel: +358 10 4261

Sverige

Orion Corporation
Tel: +358 10 4261

United Kingdom (Northern Ireland)

Orion Corporation
Tel: +358 10 4261

This leaflet was last revised in

Other sources of information

Detailed information on this medicine is available on the European Medicines Agency website:
<http://www.ema.europa.eu>.