

recepia.sk@gsk.com

Italia

GlaxoSmithKline S.p.A.
Tel: + 39 04 59 21 81 11

Suomi/Finland

GlaxoSmithKline Oy
Puh/Tel: + 358 10 30 30 30
Finland.tuoteinfo@gsk.com

Κύπρος

GlaxoSmithKline (Cyprus) Ltd
Τηλ: + 357 22 39 70 00

Sverige

GlaxoSmithKline AB
Tel: + 46 (0)8 638 93 00
info.produkt@gsk.com

Latvija

GlaxoSmithKline Latvia SIA
Tel: + 371 67312687
lv-epasts@gsk.com

United Kingdom

GlaxoSmithKline UK
Tel: + 44 (0)808 100 9997
customercontactuk@gsk.com

Lietuva

GlaxoSmithKline Lietuva UAB
Tel. +370 5 264 90 00
info.lt@gsk.com

This leaflet was last approved in

Detailed information on this medicine is available on the European Medicines Agency (EMA) web site: <http://www.ema.europa.eu/>.

The following information is intended for medical or healthcare professionals only:

Upon storage, a white deposit and clear supernatant can be observed. This does not constitute a sign of deterioration.

The syringe should be well shaken in order to obtain a homogeneous turbid white suspension.

The diphtheria, tetanus, acellular pertussis, hepatitis B, inactivated poliomyelitis (DTPa-HBV-IPV) suspension should be inspected visually for any foreign particulate matter and/or variation of physical aspect. In the event of either being observed, discard the container.