

ANNEX I
SUMMARY OF PRODUCT CHARACTERISTICS

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 150 mg film-coated tablets
Ecansya 300 mg film-coated tablets
Ecansya 500 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Ecansya 150 mg film-coated tablets

Each film-coated tablet contains 150 mg capecitabine.

Excipient with known effect

Each film-coated tablet contains 7 mg lactose.

Ecansya 300 mg film-coated tablets

Each film-coated tablet contains 300 mg capecitabine.

Excipient with known effect

Each film-coated tablet contains 15 mg lactose.

Ecansya 500 mg film-coated tablets

Each film-coated tablet contains 500 mg capecitabine.

Excipient with known effect

Each film-coated tablet contains 25 mg lactose.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Film-coated tablet (tablet).

Ecansya 150 mg film-coated tablets

The tablets are light peach colored, oblong shaped, biconvex tablets of 11.4 mm in length and 5.3 mm in width, debossed with "150" on one side and plain on other side.

Ecansya 300 mg film-coated tablets

The tablets are white to off white, oblong shaped, biconvex tablets of 14.6 mm in length and 6.7 mm in width, debossed with "300" on one side and plain on other side.

Ecansya 500 mg film-coated tablets

The tablets are peach colored, oblong shaped, biconvex tablets of 15.9 mm in length and 8.4 mm in width, debossed with "500" on one side and plain on other side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Ecansya is indicated for the treatment of:

- for the adjuvant treatment of patients following surgery of stage III (Dukes' stage C) colon cancer (see section 5.1).
- metastatic colorectal cancer (see section 5.1).

- first-line treatment of advanced gastric cancer in combination with a platinum based regimen (see section 5.1).
- in combination with docetaxel (see section 5.1) for the treatment of patients with locally advanced or metastatic breast cancer after failure of cytotoxic chemotherapy. Previous therapy should have included an anthracycline.
- as monotherapy for the treatment of patients with locally advanced or metastatic breast cancer after failure of taxanes and an anthracycline containing chemotherapy regimen or for whom further anthracycline therapy is not indicated.

4.2 Posology and method of administration

Capecitabine should only be prescribed by a qualified physician experienced in the utilisation of anti-neoplastic medicinal products. Careful monitoring during the first cycle of treatment is recommended for all patients.

Treatment should be discontinued if progressive disease or intolerable toxicity is observed. Standard and reduced dose calculations according to body surface area for starting doses of Ecansya of 1250 mg/m² and 1000 mg/m² are provided in tables 1 and 2, respectively.

Posology

Recommended posology (see section 5.1):

Monotherapy

Colon, colorectal and breast cancer

Given as monotherapy, the recommended starting dose for capecitabine in the adjuvant treatment of colon cancer, in the treatment of metastatic colorectal cancer or of locally advanced or metastatic breast cancer is 1250 mg/m² administered twice daily (morning and evening; equivalent to 2500 mg/m² total daily dose) for 14 days followed by a 7-day rest period. Adjuvant treatment in patients with stage III colon cancer is recommended for a total of 6 months.

Combination therapy

Colon, colorectal and gastric cancer

In combination treatment, the recommended starting dose of capecitabine should be reduced to 800 - 1000 mg/m² when administered twice daily for 14 days followed by a 7-day rest period, or to 625 mg/m² twice daily when administered continuously (see section 5.1). For combination with irinotecan, the recommended starting dose is 800 mg/m² when administered twice daily for 14 days followed by a 7-day rest period combined with irinotecan 200 mg/m² on day 1. The inclusion of bevacizumab in a combination regimen has no effect on the starting dose of capecitabine.

Premedication to maintain adequate hydration and anti-emesis according to the cisplatin summary of product characteristics should be started prior to cisplatin administration for patients receiving the capecitabine plus cisplatin combination. Premedication with antiemetics according to the oxaliplatin summary of product characteristics is recommended for patients receiving the capecitabine plus oxaliplatin combination. Adjuvant treatment in patients with stage III colon cancer is recommended for a duration of 6 months.

Breast cancer

In combination with docetaxel, the recommended starting dose of capecitabine in the treatment of metastatic breast cancer is 1250 mg/m² twice daily for 14 days followed by a 7-day rest period, combined with docetaxel at 75 mg/m² as a 1 hour intravenous infusion every 3 weeks. Premedication with an oral corticosteroid such as dexamethasone according to the docetaxel summary of product characteristics should be started prior to docetaxel administration for patients receiving the capecitabine plus docetaxel combination.

Ecansya dose calculations

Table 1 Standard and reduced dose calculations according to body surface area for a starting dose of capecitabine of 1250 mg/m²

	Dose level 1250 mg/m ² (twice daily)					
	Full dose 1250 mg/m ²	Number of 150 mg tablets, 300 mg tablets and/or 500 mg tablets per administration (each administration to be given morning and evening)			Reduced dose (75%) 950 mg/m ²	Reduced dose (50%) 625 mg/m ²
Body Surface Area (m ²)	Dose per administration (mg)	150 mg	300 mg	500 mg	Dose per administration (mg)	Dose per administration (mg)
≤ 1.26	1500	-	-	3	1150	800
1.27 - 1.38	1650	1	-	3	1300	800
1.39 - 1.52	1800	-	1	3	1450	950
1.53 - 1.66	2000	-	-	4	1500	1000
1.67 - 1.78	2150	1	-	4	1650	1000
1.79 - 1.92	2300	-	1	4	1800	1150
1.93 - 2.06	2500	-	-	5	1950	1300
2.07 - 2.18	2650	1	-	5	2000	1300
≥ 2.19	2800	-	1	5	2150	1450

Table 2 Standard and reduced dose calculations according to body surface area for a starting dose of capecitabine of 1000 mg/m²

	Dose level 1000 mg/m ² (twice daily)					
	Full dose 1000 mg/m ²	Number of 150 mg tablets, 300 mg tablets and/or 500 mg tablets per administration (each administration to be given morning and evening)			Reduced dose (75%) 750 mg/m ²	Reduced dose (50%) 500 mg/m ²
Body Surface Area (m ²)	Dose per administration (mg)	150 mg	300 mg	500 mg	Dose per administration (mg)	Dose per administration (mg)
≤ 1.26	1150	1	-	2	800	600
1.27 - 1.38	1300	-	1	2	1000	600
1.39 - 1.52	1450	1	1	2	1100	750
1.53 - 1.66	1600	-	2	2	1200	800
1.67 - 1.78	1750	1	2	2	1300	800
1.79 - 1.92	1800	-	1	3	1400	900
1.93 - 2.06	2000	-	-	4	1500	1000
2.07 - 2.18	2150	1	-	4	1600	1050
≥ 2.19	2300	-	1	4	1750	1100

Posology adjustments during treatment

General

Toxicity due to capecitabine administration may be managed by symptomatic treatment and/or modification of the dose (treatment interruption or dose reduction). Once the dose has been reduced, it should not be increased at a later time. For those toxicities considered by the treating physician to be unlikely to become serious or life-threatening, e.g. alopecia, altered taste, nail changes, treatment can be continued at the same dose without reduction or interruption. Patients taking capecitabine should be informed of the need to interrupt treatment immediately if moderate or severe toxicity occurs. Doses of capecitabine omitted for toxicity are not replaced. The following are the recommended dose modifications for toxicity.

Table 3 Capecitabine dose reduction schedule (3 weekly cycle or continuous treatment)

Toxicity grades*	Dose changes within a treatment cycle	Dose adjustment for next cycle/dose (% of starting dose)
• Grade 1	Maintain dose level	Maintain dose level
• Grade 2		
-1st appearance	Interrupt until resolved to grade 0-1	100%
-2nd appearance		75%
-3rd appearance		50%
-4th appearance	Discontinue treatment permanently	Not applicable
• Grade 3		
-1st appearance	Interrupt until resolved to grade 0-1	75%
-2nd appearance		50%
-3rd appearance	Discontinue treatment permanently	Not applicable
• Grade 4		
-1st appearance	Discontinue permanently <i>or</i> If physician deems it to be in the patient's best interest to continue, interrupt until resolved to grade 0-1	50%
-2nd appearance	Discontinue permanently	Not applicable

*According to the National Cancer Institute of Canada Clinical Trial Group (NCIC CTG) Common Toxicity Criteria (version 1) or the Common Terminology Criteria for Adverse Events (CTCAE) of the Cancer Therapy Evaluation Program, US National Cancer Institute, version 4.0. For hand-foot syndrome and hyperbilirubinemia, see section 4.4.

Haematology

Patients with baseline neutrophil counts of $<1.5 \times 10^9/L$ and/or thrombocyte counts of $<100 \times 10^9/L$ should not be treated with capecitabine. If unscheduled laboratory assessments during a treatment cycle show that the neutrophil count drops below $1.0 \times 10^9/L$ or that the platelet count drops below $75 \times 10^9/L$, treatment with capecitabine should be interrupted.

Dose modifications for toxicity when capecitabine is used as a 3 weekly cycle in combination with other medicinal products

Dose modifications for toxicity when capecitabine is used as a 3 weekly cycle in combination with other medicinal products should be made according to table 3 above for capecitabine and according to the appropriate summary of product characteristics for the other medicinal product(s).

At the beginning of a treatment cycle, if a treatment delay is indicated for either capecitabine or the other medicinal product(s), then administration of all therapy should be delayed until the requirements for restarting all medicinal products are met.

During a treatment cycle for those toxicities considered by the treating physician not to be related to capecitabine, capecitabine should be continued and the dose of the other medicinal product should be adjusted according to the appropriate Prescribing Information.

If the other medicinal product(s) have to be discontinued permanently, capecitabine treatment can be resumed when the requirements for restarting capecitabine are met.

This advice is applicable to all indications and to all special populations.

Dose modifications for toxicity when capecitabine is used continuously in combination with other medicinal products

Dose modifications for toxicity when capecitabine is used continuously in combination with other medicinal products should be made according to table 3 above for capecitabine and according to the appropriate summary of product characteristics for the other medicinal product(s).

Posology adjustments for special populations

Hepatic impairment

Insufficient safety and efficacy data are available in patients with hepatic impairment to provide a dose adjustment recommendation. No information is available on hepatic impairment due to cirrhosis or hepatitis.

Renal impairment

Capecitabine is contraindicated in patients with severe renal impairment (creatinine clearance below 30 ml/min [Cockcroft and Gault] at baseline). The incidence of grade 3 or 4 adverse reactions in patients with moderate renal impairment (creatinine clearance 30-50 ml/min at baseline) is increased compared to the overall population. In patients with moderate renal impairment at baseline, a dose reduction to 75% for a starting dose of 1250 mg/m² is recommended. In patients with moderate renal impairment at baseline, no dose reduction is required for a starting dose of 1000 mg/m². In patients with mild renal impairment (creatinine clearance 51-80 ml/min at baseline) no adjustment of the starting dose is recommended. Careful monitoring and prompt treatment interruption is recommended if the patient develops a grade 2, 3 or 4 adverse event during treatment and subsequent dose adjustment as outlined in table 3 above. If the calculated creatinine clearance decreases during treatment to a value below 30 ml/min, Ecansya should be discontinued. These dose adjustment recommendations for renal impairment apply both to monotherapy and combination use (see also section "Elderly" below).

Elderly

During capecitabine monotherapy, no adjustment of the starting dose is needed. However, grade 3 or 4 treatment-related adverse reactions were more frequent in patients ≥60 years of age compared to younger patients.

When capecitabine was used in combination with other medicinal products, elderly patients (≥65 years) experienced more grade 3 and grade 4 adverse drug reactions, including those leading to discontinuation, compared to younger patients. Careful monitoring of patients ≥60 years of age is advisable.

- *In combination with docetaxel:* an increased incidence of grade 3 or 4 treatment-related adverse reactions and treatment-related serious adverse reactions were observed in patients 60 years of age or more (see section 5.1). For patients 60 years of age or more, a starting dose reduction of capecitabine to 75% (950 mg/m² twice daily) is recommended. If no toxicity is observed in patients ≥60 years of age treated with a reduced capecitabine starting dose in combination with docetaxel, the dose of capecitabine may be cautiously escalated to 1250 mg/m² twice daily.

Paediatric population

There is no relevant use of capecitabine in the paediatric population in the indications colon, colorectal, gastric and breast cancer.

Method of administration

Ecansya tablets should be swallowed whole with water within 30 minutes after a meal.

Ecansya tablets should not be crushed or cut.

4.3 Contraindications

- History of severe and unexpected reactions to fluoropyrimidine therapy,
- Hypersensitivity to capecitabine or to any of the excipients listed in section 6.1 or fluorouracil,
- Known complete dihydropyrimidine dehydrogenase (DPD) deficiency (see section 4.4),
- During pregnancy and lactation,
- In patients with severe leukopenia, neutropenia, or thrombocytopenia,
- In patients with severe hepatic impairment,
- In patients with severe renal impairment (creatinine clearance below 30 ml/min),
- Recent or concomitant treatment with brivudine (see section 4.4 and 4.5 for drug-drug interaction),

- If contraindications exist to any of the medicinal products in the combination regimen, that medicinal product should not be used.

4.4 Special warnings and precautions for use

Dose limiting toxicities

Dose limiting toxicities include diarrhoea, abdominal pain, nausea, stomatitis and hand-foot syndrome (hand-foot skin reaction, palmar-plantar erythrodysesthesia). Most adverse reactions are reversible and do not require permanent discontinuation of therapy, although doses may need to be withheld or reduced.

Diarrhoea. Patients with severe diarrhoea should be carefully monitored and given fluid and electrolyte replacement if they become dehydrated. Standard antidiarrhoeal treatments (e.g. loperamide) may be used. NCIC CTC grade 2 diarrhoea is defined as an increase of 4 to 6 stools/day or nocturnal stools, grade 3 diarrhoea as an increase of 7 to 9 stools/day or incontinence and malabsorption. Grade 4 diarrhoea is an increase of ≥ 10 stools/day or grossly bloody diarrhoea or the need for parenteral support. Dose reduction should be applied as necessary (see section 4.2).

Dehydration. Dehydration should be prevented or corrected at the onset. Patients with anorexia, asthenia, nausea, vomiting or diarrhoea may rapidly become dehydrated. Dehydration may cause acute renal failure, especially in patients with pre-existing compromised renal function or when capecitabine is given concomitantly with known nephrotoxic medicinal products. Acute renal failure secondary to dehydration might be potentially fatal. If grade 2 (or higher) dehydration occurs, capecitabine treatment should be immediately interrupted and the dehydration corrected. Treatment should not be restarted until the patient is rehydrated and any precipitating causes have been corrected or controlled. Dose modifications applied should be applied for the precipitating adverse event as necessary (see section 4.2).

Hand-foot syndrome (also known as hand-foot skin reaction or palmar-plantar erythrodysesthesia or chemotherapy induced acral erythema). Grade 1 hand-foot syndrome is defined as numbness, dysesthesia/paresthesia, tingling, painless swelling or erythema of the hands and/or feet and/or discomfort which does not disrupt the patient's normal activities.

Grade 2 hand-foot syndrome is painful erythema and swelling of the hands and/or feet and/or discomfort affecting the patient's activities of daily living.

Grade 3 hand-foot syndrome is moist desquamation, ulceration, blistering and severe pain of the hands and/or feet and/or severe discomfort that causes the patient to be unable to work or perform activities of daily living. Persistent or severe hand-foot syndrome (Grade 2 and above) can eventually lead to loss of fingerprints which could impact patient identification. If grade 2 or 3 hand-foot syndrome occurs, administration of capecitabine should be interrupted until the event resolves or decreases in intensity to grade 1. Following grade 3 hand-foot syndrome, subsequent doses of capecitabine should be decreased. When capecitabine and cisplatin are used in combination, the use of vitamin B6 (pyridoxine) is not advised for symptomatic or secondary prophylactic treatment of hand-foot syndrome, because of published reports that it may decrease the efficacy of cisplatin. There is some evidence that dexpanthenol is effective for hand-foot syndrome prophylaxis in patients treated with Ecansya.

Cardiotoxicity. Cardiotoxicity has been associated with fluoropyrimidine therapy, including myocardial infarction, angina, dysrhythmias, cardiogenic shock, sudden death and electrocardiographic changes (including very rare cases of QT prolongation). These adverse reactions may be more common in patients with a prior history of coronary artery disease. Cardiac arrhythmias (including ventricular fibrillation, torsade de pointes, and bradycardia), angina pectoris, myocardial infarction, heart failure and cardiomyopathy have been reported in patients receiving capecitabine. Caution must be exercised in patients with history of significant cardiac disease, arrhythmias and angina pectoris (see section 4.8).

Hypo- or hypercalcaemia. Hypo- or hypercalcaemia has been reported during capecitabine treatment. Caution must be exercised in patients with pre-existing hypo- or hypercalcaemia (see section 4.8).

Central or peripheral nervous system disease. Caution must be exercised in patients with central or peripheral nervous system disease, e.g. brain metastasis or neuropathy (see section 4.8).

Diabetes mellitus or electrolyte disturbances. Caution must be exercised in patients with diabetes mellitus or electrolyte disturbances, as these may be aggravated during capecitabine treatment.

Coumarin-derivative anticoagulation. In an interaction study with single-dose warfarin administration, there was a significant increase in the mean AUC (+57%) of S-warfarin. These results suggest an interaction, probably due to an inhibition of the cytochrome P450 2C9 isoenzyme system by capecitabine. Patients receiving concomitant capecitabine and oral coumarin-derivative anticoagulant therapy should have their anticoagulant response (INR or prothrombin time) monitored closely and the anticoagulant dose adjusted accordingly (see section 4.5).

Brivudine. Brivudine must not be administered concomitantly with capecitabine. Fatal cases have been reported following this drug interaction. There must be at least a 4-week waiting period between end of treatment with brivudine and start of capecitabine therapy. Treatment with brivudine can be started 24 hours after the last dose of capecitabine (see section 4.3 and 4.5). In the event of accidental administration of brivudine to patients being treated with capecitabine, effective measures should be taken to reduce the toxicity of capecitabine. Immediate admission to hospital is recommended. All measures should be initiated to prevent systemic infections and dehydration.

Hepatic impairment. In the absence of safety and efficacy data in patients with hepatic impairment, capecitabine use should be carefully monitored in patients with mild to moderate liver dysfunction, regardless of the presence or absence of liver metastasis. Administration of capecitabine should be interrupted if treatment-related elevations in bilirubin of $>3.0 \times \text{ULN}$ or treatment-related elevations in hepatic aminotransferases (ALT, AST) of $>2.5 \times \text{ULN}$ occur. Treatment with capecitabine monotherapy may be resumed when bilirubin decreases to $\leq 3.0 \times \text{ULN}$ or hepatic aminotransferases decrease to $\leq 2.5 \times \text{ULN}$.

Renal impairment. The incidence of grade 3 or 4 adverse reactions in patients with moderate renal impairment (creatinine clearance 30-50 ml/min) is increased compared to the overall population (see sections 4.2 and 4.3).

Dihydropyrimidine dehydrogenase (DPD) deficiency:

DPD activity is rate limiting in the catabolism of 5-fluorouracil (see Section 5.2). Patients with DPD deficiency are therefore at increased risk of fluoropyrimidines-related toxicity, including for example stomatitis, diarrhoea, mucosal inflammation, neutropenia and neurotoxicity.

DPD-deficiency related toxicity usually occurs during the first cycle of treatment or after dose increase.

Complete DPD deficiency

Complete DPD deficiency is rare (0.01-0.5% of Caucasians). Patients with complete DPD deficiency are at high risk of life-threatening or fatal toxicity and must not be treated with Ecansya (see section 4.3).

Partial DPD deficiency

Partial DPD deficiency is estimated to affect 3-9% of the Caucasian population. Patients with partial DPD deficiency are at increased risk of severe and potentially life-threatening toxicity. A reduced starting dose should be considered to limit this toxicity. DPD deficiency should be considered as a parameter to be taken into account in conjunction with other routine measures for dose reduction. Initial dose reduction may impact the efficacy of treatment. In the absence of serious toxicity, subsequent doses may be increased with careful monitoring.

Testing for DPD deficiency

Phenotype and/or genotype testing prior to the initiation of treatment with Ecansya is recommended

despite uncertainties regarding optimal pre-treatment testing methodologies. Consideration should be given to applicable clinical guidelines.

Genotypic characterisation of DPD deficiency

Pre-treatment testing for rare mutations of the DPYD gene can identify patients with DPD deficiency.

The four DPYD variants c.1905+1G>A [also known as DPYD*2A], c.1679T>G [DPYD*13], c.2846A>T and c.1236G>A/HapB3 can cause complete absence or reduction of DPD enzymatic activity. Other rare variants may also be associated with an increased risk of severe or life-threatening toxicity.

Certain homozygous and compound heterozygous mutations in the DPYD gene locus (e.g. combinations of the four variants with at least one allele of c.1905+1G>A or c.1679T>G) are known to cause complete or near complete absence of DPD enzymatic activity.

Patients with certain heterozygous DPYD variants (including c.1905+1G>A, c.1679T>G, c.2846A>T and c.1236G>A/HapB3 variants) have increased risk of severe toxicity when treated with fluoropyrimidines.

The frequency of the heterozygous c.1905+1G>A genotype in the DPYD gene in Caucasian patients is around 1%, 1.1% for c.2846A>T, 2.6-6.3% for c.1236G>A/HapB3 variants and 0.07 to 0.1% for c.1679T>G.

Data on the frequency of the four DPYD variants in other populations than Caucasian is limited. At the present, the four DPYD variants (c.1905+1G>A, c.1679T>G, c.2846A>T and c.1236G>A/HapB3) are considered virtually absent in populations of African (-American) or Asian origin.

Phenotypic characterisation of DPD deficiency

For phenotypic characterisation of DPD deficiency, the measurement of pre-therapeutic blood levels of the endogenous DPD substrate uracil (U) in plasma is recommended.

Elevated pre-treatment uracil concentrations are associated with an increased risk of toxicity. Despite uncertainties on uracil thresholds defining complete and partial DPD deficiency, a blood uracil level ≥ 16 ng/ml and < 150 ng/ml should be considered indicative of partial DPD deficiency and associated with an increased risk for fluoropyrimidine toxicity. A blood uracil level ≥ 150 ng/ml should be considered indicative of complete DPD deficiency and associated with a risk for life-threatening or fatal fluoropyrimidine toxicity.

Ophthalmologic complications: Patients should be carefully monitored for ophthalmological complications such as keratitis and corneal disorders, especially if they have a prior history of eye disorders. Treatment of eye disorders should be initiated as clinically appropriate.

Severe skin reactions: Ecansya can induce severe skin reactions such as Stevens-Johnson syndrome and Toxic Epidermal Necrolysis. Ecansya should be permanently discontinued in patients who experience a severe skin reaction during treatment.

As this medicinal product contains lactose as an excipient, patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Ecansya tablets should not be crushed or cut. In case of exposure of either patient or caregiver to crushed or cut Ecansya tablets adverse drug reactions could occur (see Section 4.8).

4.5 Interaction with other medicinal products and other forms of interaction

Interaction studies have only been performed in adults.

Interaction with other medicinal products

Brivudine: a clinically significant interaction between brivudine and fluoropyrimidines (e.g. capecitabine, 5-fluorouracil, tegafur), resulting from the inhibition of dihydropyrimidine dehydrogenase by brivudine, has been described. This interaction, which leads to increased fluoropyrimidine toxicity, is potentially fatal. Therefore, brivudine must not be administered concomitantly with capecitabine (see section 4.3 and 4.4). There must be at least a 4-week waiting period between end of treatment with brivudine and start of capecitabine therapy. Treatment with brivudine can be started 24 hours after the last dose of capecitabine.

Cytochrome P-450 2C9 substrates: Other than warfarin, no formal interaction studies between capecitabine and other CYP2C9 substrates have been conducted. Care should be exercised when capecitabine is co-administered with 2C9 substrates (e.g., phenytoin). See also interaction with coumarin-derivative anticoagulants below, and section 4.4.

Coumarin-derivative anticoagulants: altered coagulation parameters and/or bleeding have been reported in patients taking capecitabine concomitantly with coumarin-derivative anticoagulants such as warfarin and phenprocoumon. These reactions occurred within several days and up to several months after initiating capecitabine therapy and, in a few cases, within one month after stopping capecitabine. In a clinical pharmacokinetic interaction study, after a single 20 mg dose of warfarin, capecitabine treatment increased the AUC of S-warfarin by 57% with a 91% increase in INR value. Since metabolism of R-warfarin was not affected, these results indicate that capecitabine down-regulates isozyme 2C9, but has no effect on isozymes 1A2 and 3A4. Patients taking coumarin-derivative anticoagulants concomitantly with capecitabine should be monitored regularly for alterations in their coagulation parameters (PT or INR) and the anticoagulant dose adjusted accordingly.

Phenytoin: increased phenytoin plasma concentrations resulting in symptoms of phenytoin intoxication in single cases have been reported during concomitant use of capecitabine with phenytoin. Patients taking phenytoin concomitantly with capecitabine should be regularly monitored for increased phenytoin plasma concentrations.

Folinic acid/folic acid: a combination study with capecitabine and folinic acid indicated that folinic acid has no major effect on the pharmacokinetics of capecitabine and its metabolites. However, folinic acid has an effect on the pharmacodynamics of capecitabine and its toxicity may be enhanced by folinic acid: the maximum tolerated dose (MTD) of capecitabine alone using the intermittent regimen is 3000 mg/m² per day whereas it is only 2000 mg/m² per day when capecitabine was combined with folinic acid (30 mg orally bid). The enhanced toxicity may be relevant when switching from 5-FU/LV to a capecitabine regimen. This may also be relevant with folic acid supplementation for folate deficiency due to the similarity between folinic acid and folic acid.

Antacid: the effect of an aluminium hydroxide and magnesium hydroxide-containing antacid on the pharmacokinetics of capecitabine was investigated. There was a small increase in plasma concentrations of capecitabine and one metabolite (5'-DFCR); there was no effect on the 3 major metabolites (5'-DFUR, 5-FU and FBAL).

Allopurinol: interactions with allopurinol have been observed for 5-FU; with possible decreased efficacy of 5-FU. Concomitant use of allopurinol with capecitabine should be avoided.

Interferon alpha: the MTD of capecitabine was 2000 mg/m² per day when combined with interferon alpha-2a (3 MIU/m² per day) compared to 3000 mg/m² per day when capecitabine was used alone.

Radiotherapy: the MTD of capecitabine alone using the intermittent regimen is 3000 mg/m² per day, whereas, when combined with radiotherapy for rectal cancer, the MTD of capecitabine is 2000 mg/m² per day using either a continuous schedule or given daily Monday through Friday during a 6-week course of radiotherapy.

Oxaliplatin: no clinically significant differences in exposure to capecitabine or its metabolites, free platinum or total platinum occurred when capecitabine was administered in combination with oxaliplatin or in combination with oxaliplatin and bevacizumab.

Bevacizumab: there was no clinically significant effect of bevacizumab on the pharmacokinetic parameters of capecitabine or its metabolites in the presence of oxaliplatin.

Food interaction

In all clinical trials, patients were instructed to administer capecitabine within 30 minutes after a meal. Since current safety and efficacy data are based upon administration with food, it is recommended that capecitabine be administered with food. Administration with food decreases the rate of capecitabine absorption (see section 5.2).

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception in males and females

Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with capecitabine. If the patient becomes pregnant while receiving capecitabine, the potential hazard to the foetus must be explained. An effective method of contraception should be used during treatment and for 6 months after the last dose of capecitabine.

Based on genetic toxicity findings, male patients with female partners of reproductive potential should use effective contraception during treatment and for 3 months following the last dose of capecitabine.

Pregnancy

There are no studies in pregnant women using capecitabine; however, it should be assumed that capecitabine may cause foetal harm if administered to pregnant women. In reproductive toxicity studies in animals, capecitabine administration caused embryoletality and teratogenicity. These findings are expected effects of fluoropyrimidine derivatives. Capecitabine is contraindicated during pregnancy.

Breast-feeding

It is not known whether capecitabine is excreted in human breast milk. No studies have been conducted to assess the impact of capecitabine on milk production or its presence in human breast milk. In lactating mice, considerable amounts of capecitabine and its metabolites were found in milk. As the potential for harm to the nursing infant is unknown, breast-feeding should be discontinued while receiving treatment with capecitabine and for 2 weeks after the final dose.

Fertility

There is no data on Ecansya and impact on fertility. The Ecansya pivotal studies included females of childbearing potential and males only if they agreed to use an acceptable method of birth control to avoid pregnancy for the duration of the study and for a reasonable period thereafter.

In animal studies effects on fertility were observed (see section 5.3).

4.7 Effects on ability to drive and use machines

Ecansya has minor or moderate influence on the ability to drive and use machines. Capecitabine may cause dizziness, fatigue and nausea.

4.8 Undesirable effects

Summary of the safety profile

The overall safety profile of capecitabine is based on data from over 3000 patients treated with capecitabine as monotherapy or capecitabine in combination with different chemotherapy regimens in multiple indications. The safety profiles of capecitabine monotherapy for the metastatic breast cancer, metastatic colorectal cancer and adjuvant colon cancer populations are comparable. See section 5.1 for details of major studies, including study designs and major efficacy results.

The most commonly reported and/or clinically relevant treatment-related adverse drug reactions (ADRs) were gastrointestinal disorders (especially diarrhoea, nausea, vomiting, abdominal pain, stomatitis), hand-foot syndrome (palmar-plantar erythrodysesthesia), fatigue, asthenia, anorexia, cardiotoxicity, increased renal dysfunction on those with preexisting compromised renal function, and thrombosis/embolism.

Tabulated list of adverse reactions

ADRs considered by the investigator to be possibly, probably, or remotely related to the administration of capecitabine are listed in table 4 for capecitabine given as monotherapy and in table 5 for capecitabine given in combination with different chemotherapy regimens in multiple indications. The following headings are used to rank the ADRs by frequency: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$). Within each frequency grouping, ADRs are presented in order of decreasing seriousness.

Capecitabine Monotherapy

Table 4 lists ADRs associated with the use of capecitabine monotherapy based on a pooled analysis of safety data from three major studies including over 1900 patients (studies M66001, SO14695, and SO14796). ADRs are added to the appropriate frequency grouping according to the overall incidence from the pooled analysis.

Table 4 Summary of related ADRs reported in patients treated with capecitabine monotherapy

Body System	Very Common <i>All grades</i>	Common <i>All grades</i>	Uncommon <i>Severe and/or Life-threatening (grade 3-4) or considered medically relevant</i>	Rare/Very Rare (Post-Marketing Experience)
<i>Infections and infestations</i>	-	Herpes viral infection, Nasopharyngitis, Lower respiratory tract infection	Sepsis, Urinary tract infection, Cellulitis, Tonsillitis, Pharyngitis, Oral candidiasis, Influenza, Gastroenteritis, Fungal infection, Infection, Tooth abscess	
<i>Neoplasm benign, malignant and unspecified</i>	-	-	Lipoma	
<i>Blood and lymphatic system disorders</i>	-	Neutropenia, Anaemia	Febrile neutropenia, Pancytopenia, Granulocytopenia, Thrombocytopenia, Leukopenia, Haemolytic anaemia, International Normalised Ratio (INR) increased/ Prothrombin time prolonged	
<i>Immune system disorders</i>	-	-	Hypersensitivity	Angioedema (rare)
<i>Metabolism and nutrition disorders</i>	Anorexia	Dehydration, Weight decreased	Diabetes, Hypokalaemia, Appetite disorder,	

Body System	Very Common <i>All grades</i>	Common <i>All grades</i>	Uncommon <i>Severe and/or Life-threatening (grade 3-4) or considered medically relevant</i>	Rare/Very Rare (Post-Marketing Experience)
			Malnutrition, Hypertriglyceridaemia	
<i>Psychiatric disorders</i>	-	Insomnia, Depression	Confusional state, Panic attack, Depressed mood, Libido decreased	
<i>Nervous system disorders</i>	-	Headache, Lethargy, Dizziness, Paraesthesia, Dysgeusia	Aphasia, Memory impairment, Ataxia, Syncope, Balance disorder, Sensory disorder, Neuropathy peripheral	Toxic leukoencephalopathy (very rare)
<i>Eye disorders</i>	-	Lacrimation increased, Conjunctivitis, Eye irritation	Visual acuity reduced, Diplopia	Lacrimal duct stenosis (rare), Corneal disorders (rare), keratitis (rare), punctate keratitis (rare)
<i>Ear and labyrinth disorders</i>	-	-	Vertigo, Ear pain	
<i>Cardiac disorders</i>	-	-	Angina unstable, Angina pectoris, Myocardial ischaemia/infarction, Atrial fibrillation, Arrhythmia, Tachycardia, Sinus tachycardia, Palpitations	Ventricular fibrillation (rare), QT prolongation (rare), Torsade de pointes (rare), Bradycardia (rare), Vasospasm (rare)
<i>Vascular disorders</i>	-	Thrombophlebitis	Deep vein thrombosis, Hypertension, Petechiae, Hypotension, Hot flush, Peripheral coldness	
<i>Respiratory, thoracic and mediastinal disorders</i>	-	Dyspnoea, Epistaxis, Cough, Rhinorrhoea	Pulmonary embolism, Pneumothorax, Haemoptysis, Asthma, Dyspnoea exertional	
<i>Gastrointestinal disorders</i>	Diarrhoea, Vomiting, Nausea, Stomatitis, Abdominal pain	Gastrointestinal haemorrhage, Constipation, Upper abdominal pain, Dyspepsia, Flatulence, Dry mouth	Intestinal obstruction, Ascites, Enteritis, Gastritis, Dysphagia, Abdominal pain lower, Oesophagitis, Abdominal discomfort, Gastrooesophageal reflux disease, Colitis,	

Body System	Very Common <i>All grades</i>	Common <i>All grades</i>	Uncommon <i>Severe and/or Life-threatening (grade 3-4) or considered medically relevant</i>	Rare/Very Rare (Post-Marketing Experience)
			Blood in stool	
<i>Hepatobiliary disorders</i>	-	Hyperbilirubinemia, Liver function test abnormalities	Jaundice	Hepatic failure (rare), Cholestatic hepatitis (rare)
<i>Skin and subcutaneous tissue disorders</i>	Palmar-plantar erythrodysesthesia syndrome**	Rash, Alopecia, Erythema, Dry skin, Pruritus, Skin hyper-pigmentation, Rash macular, Skin desquamation, Dermatitis, Pigmentation disorder, Nail disorder	Blister, Skin ulcer, Rash, Urticaria, Photosensitivity reaction, Palmar erythema, Swelling face, Purpura, Radiation recall syndrome	Cutaneous lupus erythematosus (rare), Severe skin reactions such as Stevens-Johnson Syndrome and toxic Epidermal Necrolysis (very rare) (see section 4.4.)
<i>Musculoskeletal and connective tissue disorders</i>	-	Pain in extremity, Back pain, Arthralgia	Joint swelling, Bone pain, Facial pain, Musculoskeletal stiffness, Muscular weakness	
<i>Renal and urinary disorders</i>	-	-	Hydronephrosis, Urinary incontinence, Haematuria, Nocturia, Blood creatinine increased	
<i>Reproductive system and breast disorders</i>	-	-	Vaginal haemorrhage	
<i>General disorders and administration site conditions</i>	Fatigue, Asthenia	Pyrexia, Oedema peripheral, Malaise, Chest pain	Oedema, Chills, Influenza like illness, Rigors, Body temperature increased	

** Based on the post-marketing experience, persistent or severe palmar-plantar erythrodysesthesia syndrome can eventually lead to loss of fingerprints (see section 4.4.)

Capecitabine in combination therapy

Table 5 lists ADRs associated with the use of capecitabine in combination with different chemotherapy regimens in multiple indications based on safety data from over 3000 patients. ADRs are added to the appropriate frequency grouping (Very common or Common) according to the highest incidence seen in any of the major clinical trials and are only added when they were seen **in addition to** those seen with capecitabine monotherapy or seen **at a higher frequency grouping** compared to capecitabine monotherapy (see table 4). Uncommon ADRs reported for capecitabine in combination therapy are consistent with the ADRs reported for capecitabine monotherapy or reported for monotherapy with the combination medicinal product (in literature and/or respective summary of product characteristics).

Some of the ADRs are reactions commonly seen with the combination medicinal product (e.g. peripheral sensory neuropathy with docetaxel or oxaliplatin, hypertension seen with bevacizumab); however an exacerbation by capecitabine therapy cannot be excluded.

Table 5 Summary of related ADRs reported in patients treated with capecitabine in combination treatment **in addition** to those seen with capecitabine monotherapy or seen at a **higher frequency grouping** compared to capecitabine monotherapy

Body System	Very Common <i>All grades</i>	Common <i>All grades</i>	Rare/Very Rare (Post-Marketing Experience)
<i>Infections and infestations</i>	-	Herpes zoster, Urinary tract infection, Oral candidiasis, Upper respiratory tract infection, Rhinitis, Influenza, +Infection, Oral herpes	
<i>Blood and lymphatic system disorders</i>	+Neutropenia, +Leucopenia, +Anaemia, +Neutropenic fever, Thrombocytopenia	Bone marrow depression, +Febrile Neutropenia	
<i>Immune system disorders</i>	-	Hypersensitivity	
<i>Metabolism and nutrition disorders</i>	Appetite decreased	Hypokalaemia, Hyponatraemia, Hypomagnesaemia, Hypocalcaemia, Hyperglycaemia	
<i>Psychiatric disorders</i>	-	Sleep disorder, Anxiety	
<i>Nervous system disorders</i>	Paraesthesia, Dysaesthesia, Peripheral neuropathy, Peripheral sensory neuropathy, Dysgeusia, Headache	Neurotoxicity, Tremor, Neuralgia, Hypersensitivity reaction, Hypoaesthesia	
<i>Eye disorders</i>	Lacrimation increased	Visual disorders, Dry eye, Eye pain, Visual impairment, Vision blurred	
<i>Ear and labyrinth disorders</i>	-	Tinnitus, Hypoacusis	
<i>Cardiac disorders</i>	-	Atrial fibrillation, Cardiac ischaemia/infarction	
<i>Vascular disorders</i>	Lower limb oedema, Hypertension, +Embolism and thrombosis	Flushing, Hypotension, Hypertensive crisis, Hot flush, Phlebitis	
<i>Respiratory, thoracic and mediastinal system disorders</i>	Sore throat, Dysaesthesia pharynx	Hiccups, Pharyngolaryngeal pain, Dysphonia	
<i>Gastrointestinal disorders</i>	Constipation, Dyspepsia	Upper gastrointestinal haemorrhage, Mouth ulceration, Gastritis, Abdominal distension, Gastroesophageal reflux disease, Oral pain, Dysphagia, Rectal haemorrhage, Abdominal pain lower, Oral dysaesthesia, Paraesthesia oral, Hypoaesthesia oral, Abdominal discomfort	
<i>Hepatobiliary disorders</i>	-	Hepatic function abnormal	
<i>Skin and subcutaneous tissue disorders</i>	Alopecia, Nail disorder	Hyperhidrosis, Rash erythematous, Urticaria, Night sweats	

Body System	Very Common <i>All grades</i>	Common <i>All grades</i>	Rare/Very Rare (Post-Marketing Experience)
<i>Musculoskeletal and connective tissue disorders</i>	Myalgia, Arthralgia, Pain in extremity	Pain in jaw, Muscle spasms, Trismus, Muscular weakness	
<i>Renal and urinary disorders</i>	-	Haematuria, Proteinuria, Creatinine renal clearance decreased, Dysuria	Acute renal failure secondary to dehydration (rare)
<i>General disorders and administration site conditions</i>	Pyrexia, Weakness, ⁺ Lethargy, Temperature intolerance	Mucosal inflammation, Pain in limb, Pain, Chills, Chest pain, Influenza-like illness, ⁺ Fever, Infusion related reaction, Injection site reaction, Infusion site pain, Injection site pain	
<i>Injury, poisoning and procedural complications</i>	-	Contusion	

⁺ For each term, the frequency count was based on ADRs of all grades. For terms marked with a “+”, the frequency count was based on grade 3-4 ADRs. ADRs are added according to the highest incidence seen in any of the major combination trials.

Description of selected adverse reactions

Hand-foot syndrome (see section 4.4):

For the capecitabine dose of 1250 mg/m² twice daily on days 1 to 14 every 3 weeks, a frequency of 53% to 60% of all-grades HFS was observed in capecitabine monotherapy trials (comprising studies in adjuvant therapy in colon cancer, treatment of metastatic colorectal cancer, and treatment of breast cancer) and a frequency of 63% was observed in the capecitabine/docetaxel arm for the treatment of metastatic breast cancer. For the capecitabine dose of 1000 mg/m² twice daily on days 1 to 14 every 3 weeks, a frequency of 22% to 30% of all-grade HFS was observed in capecitabine combination therapy.

A meta-analysis of 14 clinical trials with data from over 4700 patients treated with capecitabine monotherapy or capecitabine in combination with different chemotherapy regimens in multiple indications (colon, colorectal, gastric and breast cancer) showed that HFS (all grades) occurred in 2066 (43%) patients after a median time of 239 [95% CI 201, 288] days after starting treatment with capecitabine. In all studies combined, the following covariates were statistically significantly associated with an increased risk of developing HFS: increasing capecitabine starting dose (gram), decreasing cumulative capecitabine dose (0.1*kg), increasing relative dose intensity in the first six weeks, increasing duration of study treatment (weeks), increasing age (by 10 year increments), female gender, and good ECOG performance status at baseline (0 versus ≥1).

Diarrhoea (see section 4.4):

Capecitabine can induce the occurrence of diarrhoea, which has been observed in up to 50% of patients.

The results of a meta-analysis of 14 clinical trials with data from over 4700 patients treated with capecitabine showed that in all studies combined, the following covariates were statistically significantly associated with an increased risk of developing diarrhoea: increasing capecitabine starting dose (gram), increasing duration of study treatment (weeks), increasing age (by 10 year increments), and female gender. The following covariates were statistically significantly associated with a decreased risk of developing diarrhoea: increasing cumulative capecitabine dose (0.1*kg) and

increasing relative dose intensity in the first six weeks.

Cardiotoxicity (see section 4.4):

In addition to the ADRs described in tables 4 and 5, the following ADRs with an incidence of less than 0.1% were associated with the use of capecitabine monotherapy based on a pooled analysis from clinical safety data from 7 clinical trials including 949 patients (2 phase III and 5 phase II clinical trials in metastatic colorectal cancer and metastatic breast cancer): cardiomyopathy, cardiac failure, sudden death, and ventricular extrasystoles.

Encephalopathy:

In addition to the ADRs described in tables 4 and 5, and based on the above pooled analysis from clinical safety data from 7 clinical trials, encephalopathy was also associated with the use of capecitabine monotherapy with an incidence of less than 0.1%.

Exposure to crushed or cut capecitabine tablets:

In the instance of exposure to crushed or cut capecitabine tablets, the following adverse drug reactions have been reported: eye irritation, eye swelling, skin rash, headache, paresthesia, diarrhea, nausea, gastric irritation, and vomiting.

Special populations

Elderly (see section 4.2):

An analysis of safety data in patients ≥ 60 years of age treated with capecitabine monotherapy and an analysis of patients treated with capecitabine plus docetaxel combination therapy showed an increase in the incidence of treatment-related grade 3 and 4 adverse reactions and treatment-related serious adverse reactions compared to patients < 60 years of age. Patients ≥ 60 years of age treated with capecitabine plus docetaxel also had more early withdrawals from treatment due to adverse reactions compared to patients < 60 years of age.

The results of a meta-analysis of 14 clinical trials with data from over 4700 patients treated with capecitabine showed that in all studies combined, increasing age (by 10 year increments) was statistically significantly associated with an increased risk of developing HFS and diarrhoea and with a decreased risk of developing neutropenia.

Gender

The results of a meta-analysis of 14 clinical trials with data from over 4700 patients treated with capecitabine showed that in all studies combined, female gender was statistically significantly associated with an increased risk of developing HFS and diarrhoea and with a decreased risk of developing neutropenia.

Renal impairment (see section 4.2, 4.4, and 5.2):

An analysis of safety data in patients treated with capecitabine monotherapy (colorectal cancer) with baseline renal impairment showed an increase in the incidence of treatment-related grade 3 and 4 adverse reactions compared to patients with normal renal function (36% in patients without renal impairment n=268, vs. 41% in mild n=257 and 54% in moderate n=59, respectively) (see section 5.2). Patients with moderately impaired renal function show an increased rate of dose reduction (44%) vs. 33% and 32% in patients with no or mild renal impairment and an increase in early withdrawals from treatment (21% withdrawals during the first two cycles) vs. 5% and 8% in patients with no or mild renal impairment.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

4.9 Overdose

Symptoms

The manifestations of acute overdose include nausea, vomiting, diarrhoea, mucositis, gastrointestinal irritation and bleeding, and bone marrow depression.

Management

Medical management of overdose should include customary therapeutic and supportive medical interventions aimed at correcting the presenting clinical manifestations and preventing their possible complications.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: antineoplastic agents, antimetabolites, ATC code: L01BC06.

Capecitabine is a non-cytotoxic fluoropyrimidine carbamate, which functions as an orally administered precursor of the cytotoxic moiety 5-fluorouracil (5-FU). Capecitabine is activated via several enzymatic steps (see section 5.2). The enzyme involved in the final conversion to 5-FU, thymidine phosphorylase (ThyPase), is found in tumour tissues, but also in normal tissues, albeit usually at lower levels. In human cancer xenograft models capecitabine demonstrated a synergistic effect in combination with docetaxel, which may be related to the upregulation of thymidine phosphorylase by docetaxel.

There is evidence that the metabolism of 5-FU in the anabolic pathway blocks the methylation reaction of deoxyuridylic acid to thymidylic acid, thereby interfering with the synthesis of deoxyribonucleic acid (DNA). The incorporation of 5-FU also leads to inhibition of RNA and protein synthesis. Since DNA and RNA are essential for cell division and growth, the effect of 5-FU may be to create a thymidine deficiency that provokes unbalanced growth and death of a cell. The effects of DNA and RNA deprivation are most marked on those cells which proliferate more rapidly and which metabolise 5-FU at a more rapid rate.

Colon and colorectal cancer:

Monotherapy with capecitabine in adjuvant colon cancer

Data from one multicentre, randomised, controlled phase III clinical trial in patients with stage III (Dukes' C) colon cancer supports the use of capecitabine for the adjuvant treatment of patients with colon cancer (XACT Study; M66001). In this trial, 1987 patients were randomised to treatment with capecitabine (1250 mg/m² twice daily for 2 weeks followed by a 1-week rest period and given as 3-week cycles for 24 weeks) or 5-FU and leucovorin (Mayo Clinic regimen: 20 mg/m² leucovorin intravenous followed by 425 mg/m² intravenous bolus 5-FU, on days 1 to 5, every 28 days for 24 weeks). Capecitabine was at least equivalent to intravenous 5-FU/LV in disease-free survival in per protocol population (hazard ratio 0.92; 95% CI 0.80-1.06). In the all-randomised population, tests for difference of capecitabine vs 5-FU/LV in disease-free and overall survival showed hazard ratios of 0.88 (95% CI 0.77 – 1.01; p = 0.068) and 0.86 (95% CI 0.74 – 1.01; p = 0.060), respectively. The median follow up at the time of the analysis was 6.9 years. In a preplanned multivariate Cox analysis, superiority of capecitabine compared with bolus 5-FU/LV was demonstrated. The following factors were pre-specified in the statistical analysis plan for inclusion in the model: age, time from surgery to randomization, gender, CEA levels at baseline, lymph nodes at baseline, and country. In the all-randomised population, capecitabine was shown to be superior to 5FU/LV for disease-free survival (hazard ratio 0.849; 95% CI 0.739 - 0.976; p = 0.0212), as well as for overall survival (hazard ratio 0.828; 95% CI 0.705 - 0.971; p = 0.0203).

Combination therapy in adjuvant colon cancer

Data from one multicentre, randomised, controlled phase 3 clinical trial in patients with stage III (Dukes' C) colon cancer supports the use of capecitabine in combination with oxaliplatin (XELOX)

for the adjuvant treatment of patients with colon cancer (NO16968 study). In this trial, 944 patients were randomised to 3-week cycles for 24 weeks with capecitabine (1000 mg/m² twice daily for 2 weeks followed by a 1-week rest period) in combination with oxaliplatin (130 mg/m² intravenous infusion over 2-hours on day 1 every 3 weeks); 942 patients were randomised to bolus 5-FU and leucovorin. In the primary analysis for DFS in the ITT population, XELOX was shown to be significantly superior to 5-FU/LV (HR=0.80, 95% CI=[0.69; 0.93]; p=0.0045). The 3 year DFS rate was 71% for XELOX versus 67% for 5-FU/LV. The analysis for the secondary endpoint of RFS supports these results with a HR of 0.78 (95% CI=[0.67; 0.92]; p=0.0024) for XELOX vs. 5-FU/LV. XELOX showed a trend towards superior OS with a HR of 0.87 (95% CI=[0.72; 1.05]; p=0.1486) which translates into a 13% reduction in risk of death. The 5 year OS rate was 78% for XELOX versus 74% for 5-FU/LV. The efficacy data is based on a median observation time of 59 months for OS and 57 months for DFS. The rate of withdrawal due to adverse events was higher in the XELOX combination therapy arm (21%) as compared with that of the 5-FU/LV monotherapy arm (9%) in the ITT population.

Monotherapy with capecitabine in metastatic colorectal cancer

Data from two identically-designed, multicentre, randomised, controlled phase III clinical trials (SO14695; SO14796) support the use of capecitabine for first line treatment of metastatic colorectal cancer. In these trials, 603 patients were randomised to treatment with capecitabine (1250 mg/m² twice daily for 2 weeks followed by a 1-week rest period and given as 3-week cycles). 604 patients were randomised to treatment with 5-FU and leucovorin (Mayo regimen: 20 mg/m² leucovorin intravenous followed by 425 mg/m² intravenous bolus 5-FU, on days 1 to 5, every 28 days). The overall objective response rates in the all-randomised population (investigator assessment) were 25.7% (capecitabine) vs. 16.7% (Mayo regimen); p <0.0002. The median time to progression was 140 days (capecitabine) vs. 144 days (Mayo regimen). Median survival was 392 days (capecitabine) vs. 391 days (Mayo regimen). Currently, no comparative data are available on capecitabine monotherapy in colorectal cancer in comparison with first line combination regimens.

Combination therapy in first-line treatment of metastatic colorectal cancer

Data from a multicentre, randomised, controlled phase III clinical study (NO16966) support the use of capecitabine in combination with oxaliplatin or in combination with oxaliplatin and bevacizumab for the first-line treatment of metastatic colorectal cancer. The study contained two parts: an initial 2-arm part in which 634 patients were randomised to two different treatment groups, including XELOX or FOLFOX-4, and a subsequent 2x2 factorial part in which 1401 patients were randomised to four different treatment groups, including XELOX plus placebo, FOLFOX-4 plus placebo, XELOX plus bevacizumab, and FOLFOX-4 plus bevacizumab. See table 6 for treatment regimens.

Table 6 Treatment regimens in study NO16966 (mCRC)

	Treatment	Starting Dose	Schedule
FOLFOX-4 or FOLFOX-4 + Bevacizumab	Oxaliplatin	85 mg/m ² intravenous 2 hr	Oxaliplatin on Day 1, every 2 weeks Leucovorin on Days 1 and 2, every 2 weeks 5-fluorouracil intravenous bolus/infusion, each on Days 1 and 2, every 2 weeks
	Leucovorin	200 mg/m ² intravenous 2 hr	
	5-Fluorouracil	400 mg/m ² intravenous bolus, followed by 600 mg/ m ² intravenous 22 hr	
	Placebo or Bevacizumab	5 mg/kg intravenous 30-90 mins	Day 1, prior to FOLFOX-4, every 2 weeks
XELOX or XELOX+ Bevacizumab	Oxaliplatin	130 mg/m ² intravenous 2 hr	Oxaliplatin on Day 1, every 3 weeks Capecitabine oral twice daily for 2 weeks (followed by 1 week off- treatment)
	Capecitabine	1000 mg/m ² oral twice daily	

	Treatment	Starting Dose	Schedule
	Placebo or Bevacizumab	7.5 mg/kg intravenous 30-90 mins	Day 1, prior to XELOX, every 3 weeks
5-Fluorouracil: intravenous bolus injection immediately after leucovorin			

Non-inferiority of the XELOX-containing arms compared with the FOLFOX-4-containing arms in the overall comparison was demonstrated in terms of progression-free survival in the eligible patient population and the intent-to-treat population (see table 7). The results indicate that XELOX is equivalent to FOLFOX-4 in terms of overall survival (see table 7). A comparison of XELOX plus bevacizumab versus FOLFOX-4 plus bevacizumab was a pre-specified exploratory analysis. In this treatment subgroup comparison, XELOX plus bevacizumab was similar compared to FOLFOX-4 plus bevacizumab in terms of progression-free survival (hazard ratio 1.01; 97.5% CI 0.84 - 1.22). The median follow up at the time of the primary analyses in the intent-to-treat population was 1.5 years; data from analyses following an additional 1 year of follow up are also included in table 7. However, the on-treatment PFS analysis did not confirm the results of the general PFS and OS analysis: the hazard ratio of XELOX versus FOLFOX-4 was 1.24 with 97.5% CI 1.07 - 1.44. Although sensitivity analyses show that differences in regimen schedules and timing of tumor assessments impact the on-treatment PFS analysis, a full explanation for this result has not been found.

Table 7 Key efficacy results for the non-inferiority analysis of Study NO16966

PRIMARY ANALYSIS			
	XELOX/XELOX+P/ XELOX+BV (EPP*: N=967; ITT**: N=1017)	FOLFOX-4/FOLFOX-4+P/ FOLFOX-4+BV (EPP*: N=937; ITT**: N=1017)	
Population	Median Time to Event (Days)		HR (97.5% CI)
Parameter: Progression-free Survival			
EPP	241	259	1.05 (0.94; 1.18)
ITT	244	259	1.04 (0.93; 1.16)
Parameter: Overall Survival			
EPP	577	549	0.97 (0.84; 1.14)
ITT	581	553	0.96 (0.83; 1.12)
ADDITIONAL 1 YEAR OF FOLLOW UP			
Population	Median Time to Event (Days)		HR (97.5% CI)
Parameter: Progression-free Survival			
EPP	242	259	1.02 (0.92; 1.14)
ITT	244	259	1.01 (0.91; 1.12)
Parameter: Overall Survival			
EPP	600	594	1.00 (0.88; 1.13)
ITT	602	596	0.99 (0.88; 1.12)

*EPP=eligible patient population; **ITT=intent-to-treat population

In a randomised, controlled phase III study (CAIRO), the effect of using capecitabine at a starting dose of 1000 mg/m² for 2 weeks every 3 weeks in combination with irinotecan for the first-line treatment of patients with metastatic colorectal cancer was studied. 820 patients were randomised to receive either sequential treatment (n=410) or combination treatment (n=410). Sequential treatment consisted of first-line capecitabine (1250 mg/m² twice daily for 14 days), second-line irinotecan (350 mg/m² on day 1), and third-line combination of capecitabine (1000 mg/m² twice daily for 14 days) with oxaliplatin (130 mg/m² on day 1). Combination treatment consisted of first-line capecitabine (1000 mg/m² twice daily for 14 days) combined with irinotecan (250 mg/m² on day 1) (XELIRI) and second-line capecitabine (1000 mg/m² twice daily for 14 days) plus oxaliplatin (130 mg/m² on day 1). All treatment cycles were administered at intervals of 3 weeks. In first-line treatment the median progression-free survival in the intent-to-treat population was 5.8 months (95%CI 5.1 - 6.2 months)

for capecitabine monotherapy and 7.8 months (95%CI 7.0 - 8.3 months; $p=0.0002$) for XELIRI. However this was associated with an increased incidence of gastrointestinal toxicity and neutropenia during first-line treatment with XELIRI (26% and 11% for XELIRI and first line capecitabine respectively).

The XELIRI has been compared with 5-FU + irinotecan (FOLFIRI) in three randomised studies in patients with metastatic colorectal cancer. The XELIRI regimens included capecitabine 1000 mg/m² twice daily on days 1 to 14 of a three-week cycle combined with irinotecan 250 mg/m² on day 1. In the largest study (BICC-C), patients were randomised to receive either open label FOLFIRI (n=144), bolus 5-FU (mIFL) (n=145) or XELIRI (n=141) and were additionally randomised to receive either double-blind treatment with celecoxib or placebo. Median PFS was 7.6 months for FOLFIRI, 5.9 months for mIFL ($p=0.004$ for the comparison with FOLFIRI), and 5.8 months for XELIRI ($p=0.015$). Median OS was 23.1 months for FOLFIRI, 17.6 months for mIFL ($p=0.09$), and 18.9 months for XELIRI ($p=0.27$). Patients treated with XELIRI experienced excessive gastrointestinal toxicity compared with FOLFIRI (diarrhoea 48% and 14% for XELIRI and FOLFIRI respectively).

In the EORTC study patients were randomised to receive either open label FOLFIRI (n=41) or XELIRI (n=44) with additional randomisation to either double-blind treatment with celecoxib or placebo. Median PFS and overall survival (OS) times were shorter for XELIRI versus FOLFIRI (PFS 5.9 versus 9.6 months and OS 14.8 versus 19.9 months), in addition to which excessive rates of diarrhoea were reported in patients receiving the XELIRI regimen (41% XELIRI, 5.1% FOLFIRI).

In the study published by Skof et al, patients were randomised to receive either FOLFIRI or XELIRI. Overall response rate was 49% in the XELIRI and 48% in the FOLFIRI arm ($p=0.76$). At the end of treatment, 37% of patients in the XELIRI and 26% of patients in the FOLFIRI arm were without evidence of the disease ($p=0.56$). Toxicity was similar between treatments with the exception of neutropenia reported more commonly in patients treated with FOLFIRI.

Montagnani et al used the results from the above three studies to provide an overall analysis of randomised studies comparing FOLFIRI and XELIRI treatment regimens in the treatment of mCRC. A significant reduction in the risk of progression was associated with FOLFIRI (HR, 0.76; 95%CI, 0.62-0.95; $P < 0.01$), a result partly due to poor tolerance to the XELIRI regimens used.

Data from a randomised clinical study (Souglakos et al, 2012) comparing FOLFIRI + bevacizumab with XELIRI + bevacizumab showed no significant differences in PFS or OS between treatments. Patients were randomised to receive either FOLFIRI plus bevacizumab (Arm-A, n=167) or XELIRI plus bevacizumab (Arm-B, n=166). For Arm B, the XELIRI regimen used capecitabine 1000 mg/m² twice daily for 14 days + irinotecan 250 mg/m² on day 1. Median progression-free survival (PFS) was 10.0 and 8.9 months; $p=0.64$, overall survival 25.7 and 27.5 months; $p=0.55$ and response rates 45.5 and 39.8%; $p=0.32$ for FOLFIRI-Bev and XELIRI-Bev, respectively. Patients treated with XELIRI + bevacizumab reported a significantly higher incidence of diarrhoea, febrile neutropenia and hand-foot skin reactions than patients treated with FOLFIRI + bevacizumab with significantly increased treatment delays, dose reductions and treatment discontinuations.

Data from a multicentre, randomised, controlled phase II study (AIO KRK 0604) supports the use of capecitabine at a starting dose of 800 mg/m² for 2 weeks every 3 weeks in combination with irinotecan and bevacizumab for the first-line treatment of patients with metastatic colorectal cancer. 120 Patients were randomised to a modified XELIRI regimen with capecitabine 800 mg/m² twice daily for two weeks followed by a 7-day rest period), irinotecan (200 mg/m² as a 30 minute infusion on day 1 every 3 weeks), and bevacizumab (7.5 mg/kg as a 30 to 90 minute infusion on day 1 every 3 weeks); 127 patients were randomised to treatment with capecitabine (1000 mg/m² twice daily for two weeks followed by a 7-day rest period), oxaliplatin (130 mg/m² as a 2 hour infusion on day 1 every 3 weeks), and bevacizumab (7.5 mg/kg as a 30 to 90 minute infusion on day 1 every 3 weeks). Following a mean duration of follow-up for the study population of 26.2 months, treatment responses were as shown below:

Table 8 Key efficacy results for AIO KRK study

	XELOX + bevacizumab <i>(ITT: N=127)</i>	Modified XELIRI+ bevacizumab <i>(ITT: N= 120)</i>	Hazard ratio 95% CI P value
Progression-free Survival after 6 months			
ITT	76%	84%	-
95% CI	69 - 84%	77 - 90%	
Median progression free survival			
ITT	10.4 months	12.1 months	0.93
95% CI	9.0 - 12.0	10.8 - 13.2	0.82 - 1.07 P=0.30
Median overall survival			
ITT	24.4 months	25.5 months	0.90
95% CI	19.3 - 30.7	21.0 - 31.0	0.68 - 1.19 P=0.45

Combination therapy in second-line treatment of metastatic colorectal cancer

Data from a multicentre, randomised, controlled phase III clinical study (NO16967) support the use of capecitabine in combination with oxaliplatin for the second-line treatment of metastatic colorectal cancer. In this trial, 627 patients with metastatic colorectal carcinoma who have received prior treatment with irinotecan in combination with a fluoropyrimidine regimen as first line therapy were randomised to treatment with XELOX or FOLFOX-4. For the dosing schedule of XELOX and FOLFOX-4 (without addition of placebo or bevacizumab), refer to table 6. XELOX was demonstrated to be non-inferior to FOLFOX-4 in terms of progression-free survival in the per-protocol population and intent-to-treat population (see table 9). The results indicate that XELOX is equivalent to FOLFOX-4 in terms of overall survival (see table 9). The median follow up at the time of the primary analyses in the intent-to-treat population was 2.1 years; data from analyses following an additional 6 months of follow up are also included in table 9.

Table 9 Key efficacy results for the non-inferiority analysis of Study NO16967

PRIMARY ANALYSIS			
	XELOX (PPP*: N=251; ITT**: N=313)	FOLFOX-4 (PPP*: N=252; ITT**: N=314)	
Population	Median Time to Event (Days)		HR (95% CI)
Parameter: Progression-free Survival			
PPP	154	168	1.03 (0.87; 1.24)
ITT	144	146	0.97 (0.83; 1.14)
Parameter: Overall Survival			
PPP	388	401	1.07 (0.88; 1.31)
ITT	363	382	1.03 (0.87; 1.23)
ADDITIONAL 6 MONTHS OF FOLLOW UP			
Population	Median Time to Event (Days)		HR (95% CI)
Parameter: Progression-free Survival			
PPP	154	166	1.04 (0.87; 1.24)
ITT	143	146	0.97 (0.83; 1.14)
Parameter: Overall Survival			
PPP	393	402	1.05 (0.88; 1.27)
ITT	363	382	1.02 (0.86; 1.21)

*PPP=per-protocol population; **ITT=intent-to-treat population

Advanced gastric cancer:

Data from a multicentre, randomised, controlled phase III clinical trial in patients with advanced gastric cancer supports the use of capecitabine for the first-line treatment of advanced gastric cancer

(ML17032). In this trial, 160 patients were randomised to treatment with capecitabine (1000 mg/m² twice daily for 2 weeks followed by a 7-day rest period) and cisplatin (80 mg/m² as a 2-hour infusion every 3 weeks). A total of 156 patients were randomised to treatment with 5-FU (800 mg/m² per day, continuous infusion on days 1 to 5 every 3 weeks) and cisplatin (80 mg/m² as a 2-hour infusion on day 1, every 3 weeks). Capecitabine in combination with cisplatin was non-inferior to 5-FU in combination with cisplatin in terms of progression-free survival in the per protocol analysis (hazard ratio 0.81; 95% CI 0.63 - 1.04). The median progression-free survival was 5.6 months (capecitabine + cisplatin) versus 5.0 months (5-FU + cisplatin). The hazard ratio for duration of survival (overall survival) was similar to the hazard ratio for progression-free survival (hazard ratio 0.85; 95% CI 0.64 - 1.13). The median duration of survival was 10.5 months (capecitabine + cisplatin) versus 9.3 months (5-FU + cisplatin).

Data from a randomised multicentre, phase III study comparing capecitabine to 5-FU and oxaliplatin to cisplatin in patients with advanced gastric cancer supports the use of capecitabine for the first-line treatment of advanced gastric cancer (REAL-2). In this trial, 1002 patients were randomised in a 2x2 factorial design to one of the following 4 arms:

- ECF: epirubicin (50 mg/m² as a bolus on day 1 every 3 weeks), cisplatin (60 mg/m² as a two hour infusion on day 1 every 3 weeks) and 5-FU (200 mg/m² daily given by continuous infusion via a central line).
- ECX: epirubicin (50 mg/m² as a bolus on day 1 every 3 weeks), cisplatin (60 mg/m² as a two hour infusion on day 1 every 3 weeks), and capecitabine (625 mg/m² twice daily continuously).
- EOF: epirubicin (50 mg/m² as a bolus on day 1 every 3 weeks), oxaliplatin (130 mg/m² given as a 2 hour infusion on day 1 every three weeks), and 5-FU (200 mg/m² daily given by continuous infusion via a central line).
- EOX: epirubicin (50 mg/m² as a bolus on day 1 every 3 weeks), oxaliplatin (130 mg/m² given as a 2 hour infusion on day 1 every three weeks), and capecitabine (625 mg/m² twice daily continuously).

The primary efficacy analyses in the per protocol population demonstrated non-inferiority in overall survival for capecitabine- vs 5-FU-based regimens (hazard ratio 0.86; 95% CI 0.8 - 0.99) and for oxaliplatin- vs cisplatin-based regimens (hazard ratio 0.92; 95% CI 0.80 - 1.1). The median overall survival was 10.9 months in capecitabine-based regimens and 9.6 months in 5-FU based regimens. The median overall survival was 10.0 months in cisplatin-based regimens and 10.4 months in oxaliplatin-based regimens.

Capecitabine has also been used in combination with oxaliplatin for the treatment of advanced gastric cancer. Studies with capecitabine monotherapy indicate that capecitabine has activity in advanced gastric cancer.

Colon, colorectal and advanced gastric cancer: meta-analysis

A meta-analysis of six clinical trials (studies SO14695, SO14796, M66001, NO16966, NO16967, M17032) supports capecitabine replacing 5-FU in mono- and combination treatment in gastrointestinal cancer. The pooled analysis includes 3097 patients treated with capecitabine-containing regimens and 3074 patients treated with 5-FU-containing regimens. Median overall survival time was 703 days (95% CI: 671; 745) in patients treated with capecitabine-containing regimens and 683 days (95% CI: 646; 715) in patients treated with 5-FU-containing regimens. The hazard ratio for overall survival was 0.94 (95% CI: 0.89; 1.00, p=0.0489) indicating that capecitabine-containing regimens are non-inferior to 5-FU-containing regimens.

Breast cancer

Combination therapy with capecitabine and docetaxel in locally advanced or metastatic breast cancer

Data from one multicentre, randomised, controlled phase III clinical trial support the use of capecitabine in combination with docetaxel for treatment of patients with locally advanced or metastatic breast cancer after failure of cytotoxic chemotherapy, including an anthracycline. In this trial, 255 patients were randomised to treatment with capecitabine (1250 mg/m² twice daily for 2

weeks followed by 1-week rest period and docetaxel 75 mg/m² as a 1 hour intravenous infusion every 3 weeks). 256 patients were randomised to treatment with docetaxel alone (100 mg/m² as a 1 hour intravenous infusion every 3 weeks). Survival was superior in the capecitabine + docetaxel combination arm (p=0.0126). Median survival was 442 days (capecitabine + docetaxel) vs. 352 days (docetaxel alone). The overall objective response rates in the all-randomised population (investigator assessment) were 41.6% (capecitabine + docetaxel) vs. 29.7% (docetaxel alone); p = 0.0058. Time to progressive disease was superior in the capecitabine + docetaxel combination arm (p<0.0001). The median time to progression was 186 days (capecitabine + docetaxel) vs. 128 days (docetaxel alone).

Monotherapy with capecitabine after failure of taxanes, anthracycline containing chemotherapy, and for whom anthracycline therapy is not indicated

Data from two multicentre phase II clinical trials support the use of capecitabine monotherapy for treatment of patients after failure of taxanes and an anthracycline-containing chemotherapy regimen or for whom further anthracycline therapy is not indicated. In these trials, a total of 236 patients were treated with capecitabine (1250 mg/m² twice daily for 2 weeks followed by 1-week rest period). The overall objective response rates (investigator assessment) were 20% (first trial) and 25% (second trial). The median time to progression was 93 and 98 days. Median survival was 384 and 373 days.

All indications:

A meta-analysis of 14 clinical trials with data from over 4700 patients treated with capecitabine monotherapy or capecitabine in combination with different chemotherapy regimens in multiple indications (colon, colorectal, gastric and breast cancer) showed that patients on capecitabine who developed hand-foot syndrome (HFS) had a longer overall survival compared to patients who did not develop HFS: median overall survival 1100 days (95% CI 1007;1200) vs 691 days (95% CI 638;754) with a hazard ratio of 0.61 (95% CI 0.56; 0.66).

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with the reference medicinal product containing capecitabine in all subsets of the paediatric population in adenocarcinoma of the colon and rectum, gastric adenocarcinoma and breast carcinoma (see section 4.2 for information on paediatric use).

5.2 Pharmacokinetic properties

The pharmacokinetics of capecitabine have been evaluated over a dose range of 502-3514 mg/m²/day. The parameters of capecitabine, 5'-deoxy-5-fluorocytidine (5'-DFCR) and 5'-deoxy-5-fluorouridine (5'-DFUR) measured on days 1 and 14 were similar. The AUC of 5-FU was 30%-35% higher on day 14. Capecitabine dose reduction decreases systemic exposure to 5-FU more than dose-proportionally, due to non-linear pharmacokinetics for the active metabolite.

Absorption

After oral administration, capecitabine is rapidly and extensively absorbed, followed by extensive conversion to the metabolites, 5'-DFCR and 5'-DFUR. Administration with food decreases the rate of capecitabine absorption, but only results in a minor effect on the AUC of 5'-DFUR, and on the AUC of the subsequent metabolite 5-FU. At the dose of 1250 mg/m² on day 14 with administration after food intake, the peak plasma concentrations (C_{max} in µg/ml) for capecitabine, 5'-DFCR, 5'-DFUR, 5-FU and FBAL were 4.67, 3.05, 12.1, 0.95 and 5.46 respectively. The time to peak plasma concentrations (T_{max} in hours) were 1.50, 2.00, 2.00, 2.00 and 3.34. The AUC_{0-∞} values in µg•h/ml were 7.75, 7.24, 24.6, 2.03 and 36.3.

Distribution

In vitro human plasma studies have determined that capecitabine, 5'-DFCR, 5'-DFUR and 5-FU are 54%, 10%, 62% and 10% protein bound, mainly to albumin.

Biotransformation

Capecitabine is first metabolised by hepatic carboxylesterase to 5'-DFCR, which is then converted to

5-DFUR by cytidine deaminase, principally located in the liver and tumour tissues. Further catalytic activation of 5'-DFUR then occurs by thymidine phosphorylase (ThyPase). The enzymes involved in the catalytic activation are found in tumour tissues but also in normal tissues, albeit usually at lower levels. The sequential enzymatic biotransformation of capecitabine to 5-FU leads to higher concentrations within tumour tissues. In the case of colorectal tumours, 5-FU generation appears to be in large part localised in tumour stromal cells. Following oral administration of capecitabine to patients with colorectal cancer, the ratio of 5-FU concentration in colorectal tumours to adjacent tissues was 3.2 (ranged from 0.9 to 8.0). The ratio of 5-FU concentration in tumour to plasma was 21.4 (ranged from 3.9 to 59.9, n=8) whereas the ratio in healthy tissues to plasma was 8.9 (ranged from 3.0 to 25.8, n=8). Thymidine phosphorylase activity was measured and found to be 4 times greater in primary colorectal tumour than in adjacent normal tissue. According to immunohistochemical studies, thymidine phosphorylase appears to be in large part localised in tumour stromal cells.

5-FU is further catabolised by the enzyme dihydropyrimidine dehydrogenase (DPD) to the much less toxic dihydro-5-fluorouracil (FUH₂). Dihydropyrimidinase cleaves the pyrimidine ring to yield 5-fluoro-ureidopropionic acid (FUPA). Finally, β -ureido-propionase cleaves FUPA to α -fluoro- β -alanine (FBAL) which is cleared in the urine. Dihydropyrimidine dehydrogenase (DPD) activity is the rate limiting step. Deficiency of DPD may lead to increased toxicity of capecitabine (see section 4.3 and 4.4).

Elimination

The elimination half-life ($t_{1/2}$ in hours) of capecitabine, 5'-DFUR, 5'-DFUR, 5-FU and FBAL were 0.85, 1.11, 0.66, 0.76 and 3.23 respectively. Capecitabine and its metabolites are predominantly excreted in urine; 95.5% of administered capecitabine dose is recovered in urine. Faecal excretion is minimal (2.6%). The major metabolite excreted in urine is FBAL, which represents 57% of the administered dose. About 3% of the administered dose is excreted in urine unchanged.

Combination therapy

Phase I studies evaluating the effect of capecitabine on the pharmacokinetics of either docetaxel or paclitaxel and vice versa showed no effect by capecitabine on the pharmacokinetics of docetaxel or paclitaxel (C_{max} and AUC) and no effect by docetaxel or paclitaxel on the pharmacokinetics of 5'-DFUR.

Pharmacokinetics in special populations

A population pharmacokinetic analysis was carried out after capecitabine treatment of 505 patients with colorectal cancer dosed at 1250 mg/m² twice daily. Gender, presence or absence of liver metastasis at baseline, Karnofsky Performance Status, total bilirubin, serum albumin, ASAT and ALAT had no statistically significant effect on the pharmacokinetics of 5'-DFUR, 5-FU and FBAL.

Hepatic impairment due to liver metastases.

According to a pharmacokinetic study in cancer patients with mild to moderate liver impairment due to liver metastases, the bioavailability of capecitabine and exposure to 5-FU may increase compared to patients with no liver impairment. There are no pharmacokinetic data on patients with severe hepatic impairment.

Renal impairment.

Based on a pharmacokinetic study in cancer patients with mild to severe renal impairment, there is no evidence for an effect of creatinine clearance on the pharmacokinetics of intact drug and 5-FU. Creatinine clearance was found to influence the systemic exposure to 5'-DFUR (35% increase in AUC when creatinine clearance decreases by 50%) and to FBAL (114% increase in AUC when creatinine clearance decreases by 50%). FBAL is a metabolite without antiproliferative activity.

Elderly.

Based on the population pharmacokinetic analysis, which included patients with a wide range of ages (27 to 86 years) and included 234 (46%) patients greater or equal to 65, age has no influence on the pharmacokinetics of 5'-DFUR and 5-FU. The AUC of FBAL increased with age (20% increase in age results in a 15% increase in the AUC of FBAL). This increase is likely due to a change in renal

function.

Ethnic factors.

Following oral administration of 825 mg/m² capecitabine twice daily for 14 days, Japanese patients (n=18) had about 36% lower C_{max} and 24% lower AUC for capecitabine than Caucasian patients (n=22). Japanese patients had also about 25% lower C_{max} and 34% lower AUC for FBAL than Caucasian patients. The clinical relevance of these differences is unknown. No significant differences occurred in the exposure to other metabolites (5'-DFCR, 5'-DFUR, and 5-FU).

5.3 Preclinical safety data

In repeat-dose toxicity studies, daily oral administration of capecitabine to cynomolgus monkeys and mice produced toxic effects on the gastrointestinal, lymphoid and haemopoietic systems, typical for fluoropyrimidines. These toxicities were reversible. Skin toxicity, characterised by degenerative/regressive changes, was observed with capecitabine. Capecitabine was devoid of hepatic and CNS toxicities. Cardiovascular toxicity (e.g. PR- and QT-interval prolongation) was detectable in cynomolgus monkeys after intravenous administration (100 mg/kg) but not after repeated oral dosing (1379 mg/m²/day).

A two-year mouse carcinogenicity study produced no evidence of carcinogenicity by capecitabine.

During standard fertility studies, impairment of fertility was observed in female mice receiving capecitabine; however, this effect was reversible after a drug-free period. In addition, during a 13-week study, atrophic and degenerative changes occurred in reproductive organs of male mice; however these effects were reversible after a drug-free period (see section 4.6).

In embryotoxicity and teratogenicity studies in mice, dose-related increases in foetal resorption and teratogenicity were observed. In monkeys, abortion and embryoletality were observed at high doses, but there was no evidence of teratogenicity.

Capecitabine was not mutagenic *in vitro* to bacteria (Ames test) or mammalian cells (Chinese hamster V79/HPRT gene mutation assay). However, similar to other nucleoside analogues (ie, 5-FU), capecitabine was clastogenic in human lymphocytes (*in vitro*) and a positive trend occurred in mouse bone marrow micronucleus tests (*in vivo*).

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core

Lactose
Microcrystalline cellulose (E460)
Croscarmellose sodium
Hypromellose (E-5)
Magnesium stearate

Tablet coating

- *Ecansya_150 mg film-coated tablets*
Hypromellose (6cps)
Talc
Titanium dioxide (E171)
Iron oxide red (E172)
Iron oxide yellow (E172)

- *Ecansya_300 mg film-coated tablets*
Hypromellose (6cps)

Talc
Titanium dioxide (E171)

- *Ecansya_500 mg film-coated tablets*

Hypromellose (6cps)

Talc

Titanium dioxide (E171)

Iron oxide red (E172)

Iron oxide yellow (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Aluminium/aluminium blisters

This medicinal product does not require any special storage conditions.

PVC/PVdC/aluminium blisters

Do not store above 30°C.

6.5 Nature and contents of container

Aluminium/aluminium or PVC/PVdC/aluminium blister containing 10 film-coated tablets. Each pack contains 30, 60 or 120 film-coated tablets.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

Procedures for safe handling of cytotoxic drugs should be followed.

7. MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

8. MARKETING AUTHORISATION NUMBER(S)

Ecansya 150 mg film-coated tablets

- *Aluminium/aluminium blisters*

30 film-coated tablets: EU/1/12/763/001

60 film-coated tablets: EU/1/12/763/002

120 Film-coated tablets: EU/1/12/763/003

- *PVC/PVdC/aluminium blisters*

30 film-coated tablets: EU/1/12/763/004

60 film-coated tablets: EU/1/12/763/005

120 film-coated tablets: EU/1/12/763/006

Ecansya 300 mg film-coated tablets

- *Aluminium/aluminium blisters*
30 film-coated tablets: EU/1/12/763/007
60 film-coated tablets: EU/1/12/763/008
120 film-coated tablets: EU/1/12/763/009

- *PVC/PVdC/aluminium blisters*
30 film-coated tablets: EU/1/12/763/010
60 film-coated tablets: EU/1/12/763/011
120 film-coated tablets: EU/1/12/763/012

Ecansya 500 mg film-coated tablets

- *Aluminium/aluminium blisters*
30 film-coated tablets: EU/1/12/763/013
60 film-coated tablets: EU/1/12/763/014
120 film-coated tablets: EU/1/12/763/015

- *PVC/PVdC/aluminium blisters*
30 film-coated tablets: EU/1/12/763/016
60 film-coated tablets: EU/1/12/763/017
120 film-coated tablets: EU/1/12/763/018

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 20 April 2012
Date of latest renewal: 9 December 2016

10. DATE OF REVISION OF THE TEXT

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

ANNEX II

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE

Name and address of the manufacturers responsible for batch release

Pharmacare Premium Ltd
HHF 003, Hal Far Industrial Estate Birzebbugia, BBG 3000
Malta

KRKA, d.d., Novo mesto
Šmarješka cesta 6
8501 Novo mesto
Slovenia

The printed package leaflet of the medicinal product must state the name and address of the manufacturer responsible for the release of the concerned batch.

B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT

- **Risk management plan (RMP)**

Not applicable.

ANNEX III
LABELLING AND PACKAGE LEAFLET

A. LABELLING

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 150 mg film-coated tablets
capecitabine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each film-coated tablet contains 150 mg capecitabine.

3. LIST OF EXCIPIENTS

Also contains lactose.
See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

30 film-coated tablets
60 film-coated tablets
120 film-coated tablets

5. METHOD AND ROUTE (S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING (S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

For PVC/PVdC/aluminium blisters

Do not store above 30°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

12. MARKETING AUTHORISATION NUMBER (S)

Aluminium/aluminium blisters

EU/1/12/763/001 30 film-coated tablets

EU/1/12/763/002 60 film-coated tablets

EU/1/12/763/003 120 Film-coated tablets

PVC/PVdC/aluminium blisters

EU/1/12/763/004 30 film-coated tablets

EU/1/12/763/005 60 film-coated tablets

EU/1/12/763/006 120 film-coated tablets

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Ecansya 150 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 150 mg film-coated tablets

multilingual blister:

Ecansya 150 mg tablets

capecitabine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 300 mg film-coated tablets
capecitabine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each film-coated tablet contains 300 mg capecitabine.

3. LIST OF EXCIPIENTS

Also contains lactose.
See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

30 film-coated tablets
60 film-coated tablets
120 film-coated tablets

5. METHOD AND ROUTE (S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING (S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

For PVC/PVdC/aluminium blisters

Do not store above 30°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

12. MARKETING AUTHORISATION NUMBER (S)

Aluminium/aluminium blisters

EU/1/12/763/007 30 film-coated tablets

EU/1/12/763/008 60 film-coated tablets

EU/1/12/763/009 120 film-coated tablets

PVC/PVdC/aluminium blisters

EU/1/12/763/010 30 film-coated tablets

EU/1/12/763/011 60 film-coated tablets

EU/1/12/763/012 120 film-coated tablets

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Ecansya 300 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 300 mg film-coated tablets

multilingual blister:

Ecansya 300 mg tablets

capecitabine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

PARTICULARS TO APPEAR ON THE OUTER PACKAGING

CARTON

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 500 mg film-coated tablets
capecitabine

2. STATEMENT OF ACTIVE SUBSTANCE(S)

Each film-coated tablet contains 500 mg capecitabine.

3. LIST OF EXCIPIENTS

Also contains lactose.
See leaflet for further information.

4. PHARMACEUTICAL FORM AND CONTENTS

Film-coated tablet

30 film-coated tablets
60 film-coated tablets
120 film-coated tablets

5. METHOD AND ROUTE (S) OF ADMINISTRATION

Read the package leaflet before use.

Oral use

6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN

Keep out of the sight and reach of children.

7. OTHER SPECIAL WARNING (S), IF NECESSARY

8. EXPIRY DATE

EXP

9. SPECIAL STORAGE CONDITIONS

For PVC/PVdC/aluminium blisters

Do not store above 30°C.

10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE

11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

12. MARKETING AUTHORISATION NUMBER (S)

Aluminium/aluminium blisters

EU/1/12/763/013 30 film-coated tablets

EU/1/12/763/014 60 film-coated tablets

EU/1/12/763/015 120 film-coated tablets

PVC/PVdC/aluminium blisters

EU/1/12/763/016 30 film-coated tablets

EU/1/12/763/017 60 film-coated tablets

EU/1/12/763/018 120 film-coated tablets

13. BATCH NUMBER

Lot

14. GENERAL CLASSIFICATION FOR SUPPLY

15. INSTRUCTIONS ON USE

16. INFORMATION IN BRAILLE

Ecansya 500 mg

17. UNIQUE IDENTIFIER – 2D BARCODE

2D barcode carrying the unique identifier included.

18. UNIQUE IDENTIFIER - HUMAN READABLE DATA

PC
SN
NN

MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS

BLISTER

1. NAME OF THE MEDICINAL PRODUCT

Ecansya 500 mg film-coated tablets

multilingual blister:

Ecansya 500 mg tablets

capecitabine

2. NAME OF THE MARKETING AUTHORISATION HOLDER

KRKA

3. EXPIRY DATE

EXP

4. BATCH NUMBER

Lot

5. OTHER

B. PACKAGE LEAFLET

Package leaflet: Information for the patient

Ecansya 150 mg film-coated tablets

Ecansya 300 mg film-coated tablets

Ecansya 500 mg film-coated tablets

capecitabine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What Ecansya is and what it is used for
2. What you need to know before you take Ecansya
3. How to take Ecansya
4. Possible side effects
5. How to store Ecansya
6. Contents of the pack and other information

1. What Ecansya is and what it is used for

Ecansya belongs to the group of medicines called "cytostatic medicines", which stop the growth of cancer cells. Ecansya contains capecitabine, which itself is not a cytostatic medicine. Only after being absorbed by the body is it changed into an active anti-cancer medicine (more in tumour tissue than in normal tissue).

Ecansya is used in the treatment of colon, rectal, gastric, or breast cancers. Furthermore, Ecansya is used to prevent new occurrence of colon cancer after complete removal of the tumour by surgery.

Ecansya may be used either alone or in combination with other medicines.

2. What you need to know before you take Ecansya

Do not take Ecansya

- if you are allergic to capecitabine or any of the other ingredients of this medicine (listed in section 6). You must inform your doctor if you know that you have an allergy or over-reaction to this medicine,
- if you previously have had severe reactions to fluoropyrimidine therapy (a group of anticancer medicines, such as fluorouracil),
- if you are pregnant or breast-feeding,
- if you have severely low levels of white cells or platelets in the blood (leucopenia, neutropenia or thrombocytopenia),
- if you have severe liver or kidney problems,
- if you know that you do not have any activity of the enzyme dihydropyrimidine dehydrogenase (DPD) (complete DPD deficiency),
- if you are being treated now or have been treated in the last 4 weeks with brivudine as part of herpes zoster (chickenpox or shingles) therapy.

Warnings and precautions

Talk to your doctor or pharmacist before taking Ecansya

- if you know that you have a partial deficiency in the activity of the enzyme dihydropyrimidine dehydrogenase (DPD)
- if you have a family member who has partial or complete deficiency of the enzyme dihydropyrimidine dehydrogenase (DPD)
- if you have liver or kidney diseases
- if you have or had heart problems (for example an irregular heartbeat or pains to the chest, jaw and back, brought on by physical effort and due to problems with the blood flow to the heart)
- if you have brain diseases (for example, cancer that has spread to the brain) or nerve damage (neuropathy)
- if you have calcium imbalances (seen in blood tests)
- if you have diabetes
- if you cannot keep food or water in your body because of severe nausea and vomiting
- if you have diarrhoea
- if you are or become dehydrated
- if you have imbalances of ions in your blood (electrolyte imbalances, seen in tests)
- if you have a history of eye problems as you may need extra monitoring of your eyes
- if you have a severe skin reaction.

DPD deficiency: DPD deficiency is a genetic condition that is not usually associated with health problems unless you receive certain medicines. If you have DPD deficiency and take Ecansya, you are at an increased risk of severe side effects (listed under section 4 Possible side effects). It is recommended to test you for DPD deficiency before start of treatment. If you have no activity of the enzyme you should not take Ecansya. If you have a reduced enzyme activity (partial deficiency) your doctor might prescribe a reduced dose. If you have negative test results for DPD deficiency, severe and life-threatening side effects may still occur.

Children and adolescents

Ecansya is not indicated in children and adolescents. Do not give Ecansya to children and adolescents.

Other medicines and Ecansya

Before starting treatment, tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines. This is extremely important, as taking more than one medicine at the same time can strengthen or weaken the effect of the medicines.

You must not take brivudine (an anti-viral medicine for the treatment of shingles or chickenpox) at the same time as capecitabine treatment (including during any rest periods when you are not taking any capecitabine tablets).

If you have taken brivudine you must wait for at least 4 weeks after stopping brivudine before starting to take capecitabine. See also section "Do not take Ecansya".

Also, you need to be particularly careful if you are taking any of the following:

- gout medicines (allopurinol),
- blood-thinning medicines (coumarin, warfarin),
- medicines for seizures or tremors (phenytoin),
- interferon alpha,
- radiotherapy and certain medicines used to treat cancer (folinic acid, oxaliplatin, bevacizumab, cisplatin, irinotecan),
- medicines used to treat folic acid deficiency.

Ecansya with food and drink

You should take Ecansya no later than 30 minutes after meals.

Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine. You must not take Ecansya if you

are pregnant or think you might be.

You must not breast-feed if you are taking Ecansya and for 2 weeks after the last dose.

If you are a woman who could become pregnant you should use effective contraception during treatment with Ecansya and for 6 months after the last dose.

If you are a male patient and your female partner could become pregnant, you should use effective contraception during treatment with Ecansya and for 3 months after the last dose.

Driving and using machines

Ecansya may make you feel dizzy, nauseous or tired. It is therefore possible that Ecansya could affect your ability to drive a car or operate machines.

Ecansya contains lactose

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

3. How to take Ecansya

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Capecitabine should only be prescribed by a doctor experienced in the use of anticancer medicines.

Your doctor will prescribe a dose and treatment regimen that is right for *you*. The dose of Ecansya is based on your body surface area. This is calculated from your height and weight. The usual dose for adults is 1250 mg/m² of body surface area taken two times daily (morning and evening). Two examples are provided here: A person whose body weight is 64 kg and height is 1.64 m has a body surface area of 1.7 m² and should take 4 tablets of 500 mg and 1 tablet of 150 mg two times daily. A person whose body weight is 80 kg and height is 1.80 m has a body surface area of 2.00 m² and should take 5 tablets of 500 mg two times daily.

Your doctor will tell you what dose you need to take, when to take it and for how long you need to take it.

Your doctor may want you to take a combination of *150 mg*, *300 mg* and *500 mg* tablets for each dose.

- Take the tablets **morning and evening** as prescribed by your doctor.
- Take the tablets within **30 minutes after the end of a meal** (breakfast and dinner) **and swallow whole with water. Do not crush or cut tablets. If you cannot swallow Ecansya tablets whole, tell your healthcare provider.**
- It is important that you take all your medicine as prescribed by your doctor.

Ecansya tablets are usually taken for 14 days followed by a 7 day rest period (when no tablets are taken). This 21 day period is one treatment cycle.

In combination with other medicines the usual dose for adults may be less than 1250 mg/m² of body surface area, and you may need to take the tablets over a different time period (e.g. every day, with no rest period).

If you take more Ecansya than you should

If you take more Ecansya than you should, contact your doctor as soon as possible before taking the next dose.

You might get the following side effects if you take a lot more capecitabine than you should: feeling or being sick, diarrhoea, inflammation or ulceration of the gut or mouth, pain or bleeding from the intestine or stomach, or bone marrow depression (reduction in certain kinds of blood cells). Tell your doctor immediately if you experience any of these symptoms.

If you forget to take Ecansya

Do not take the missed dose at all. Do not take a double dose to make up for a forgotten dose. Instead, continue your regular dosing schedule and check with your doctor.

If you stop taking Ecansya

There are no side-effects caused by stopping treatment with capecitabine. In case you are using coumarin anticoagulants (containing e.g. phenprocoumon), stopping capecitabine might require that your doctor adjusts your anticoagulant dose.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

STOP taking Ecansya immediately and contact your doctor if any of these symptoms occur:

- **Diarrhoea:** if you have an increase of 4 or more bowel movements compared to your normal bowel movements each day or any diarrhoea at night.
- **Vomiting:** if you vomit more than once in a 24-hour time period.
- **Nausea:** if you lose your appetite, and the amount of food you eat each day is much less than usual.
- **Stomatitis:** if you have pain, redness, swelling or sores in your mouth and/or throat.
- **Hand-and-foot skin-reaction:** if you have pain, swelling, redness or tingling of hands and/or feet.
- **Fever:** if you have a temperature of 38°C or greater.
- **Infection:** if you experience signs of infection caused by bacteria or virus, or other organisms.
- **Chest pain:** if you experience pain localised to the centre of the chest, especially if it occurs during exercise.
- **Steven-Johnson syndrome:** if you experience painful red or purplish rash that spreads and blisters and/or other lesions begin to appear in the mucous membrane (e.g. mouth and lips), in particular if you had before light sensitivity, infections of the respiratory system (e.g. bronchitis) and/or fever.
- **DPD deficiency:** if you have a known DPD deficiency, you are at an increased risk of acute early-onset of toxicity and severe, life-threatening, or fatal adverse reactions caused by Ecansya (e.g. stomatitis, mucosal inflammation, diarrhoea, neutropenia, and neurotoxicity).
- **Angioedema:** Seek medical attention straight away if you notice any of the following symptoms – you may need urgent medical treatment: swelling mainly of the face, lips, tongue or throat which makes it difficult to swallow or breathe, itching and rashes. This could be a sign of angioedema.

If caught early, these side effects usually improve within 2 to 3 days after treatment discontinuation. If these side effects continue, however, contact your doctor immediately. Your doctor may instruct you to restart treatment at a lower dose.

If severe stomatitis (sores in your mouth and/or throat), mucosal inflammation, diarrhoea, neutropenia (increased risk for infections), or neurotoxicity occurs during the first cycle of treatment a DPD deficiency may be involved (please see Section 2: Warning and precautions).

Hand and foot skin reaction can lead to loss of fingerprint, which could impact your identification by fingerprint scan.

In addition to the above, when Ecansya is used alone, very common side effects, which may affect more than 1 in 10 people are:

- abdominal pain
- rash, dry or itchy skin
- tiredness
- loss of appetite (anorexia)

These side effects can become severe; therefore, it is important that you **always contact your doctor immediately** when you start to experience a side effect. Your doctor may instruct you to decrease the dose and/or temporarily discontinue treatment with Ecansya. This will help reduce the likelihood that the side effect continues or becomes severe.

Other side effects are:

Common side effects (may affect up to 1 in 10 people) include:

- decreases in the number of white blood cells or red blood cells (seen in tests)
- dehydration, weight loss
- sleeplessness (insomnia), depression
- headache, sleepiness, dizziness, abnormal sensation in the skin (numbness or tingling sensation), taste changes
- eye irritation, increased tears, eye redness (conjunctivitis)
- inflammation of the veins (thrombophlebitis)
- shortness of breath, nose bleeds, cough, runny nose
- cold sores or other herpes infections
- infections of the lungs or respiratory system (e.g. pneumonia or bronchitis)
- bleeding from the gut, constipation, pain in upper abdomen, indigestion, excess wind, dry mouth
- skin rash, hair loss (alopecia), skin reddening, dry skin, itching (pruritus), skin discolouration, skin loss, skin inflammation, nail disorder
- pain in the joints or in the limbs (extremities), chest or back
- fever, swelling in the limbs, feeling ill
- problems with liver function (seen in blood tests) and increased blood bilirubin (excreted by the liver)

Uncommon side-effects (may affect up to 1 in 100 people) include:

- blood infection, urinary tract infection, infection of the skin, infections in the nose and throat, fungal infections (including those of the mouth), influenza, gastroenteritis, tooth abscess
- lumps under the skin (lipoma)
- decreases in blood cells including platelets, thinning of blood (seen in tests)
- allergy
- diabetes, decrease in blood potassium, malnutrition, increased blood triglycerides
- confusional state, panic attacks, depressed mood, decreased libido
- difficulty speaking, impaired memory, loss of movement coordination, balance disorder, fainting, nerve damage (neuropathy) and problems with sensation
- blurred or double vision
- vertigo, ear pain
- irregular heartbeat and palpitations (arrhythmias), chest pain and heart attack (infarction),
- blood clots in the deep veins, high or low blood pressure, hot flushes, cold limbs (extremities), purple spots on the skin
- blood clots in the veins in the lung (pulmonary embolism), collapsed lung, coughing up blood, asthma, shortness of breath on exertion
- bowel obstruction, collection of fluid in the abdomen, inflammation of the small or large intestine, the stomach or the oesophagus, pain in the lower abdomen, abdominal discomfort, heartburn (reflux of food from the stomach), blood in the stool
- jaundice (yellowing of skin and eyes)
- skin ulcer and blister, reaction of the skin with sunlight, reddening of palms, swelling or pain of the face
- joint swelling or stiffness, bone pain, muscle weakness or stiffness
- fluid collection in the kidneys, increased frequency of urination during the night, incontinence, blood in the urine, increase in blood creatinine (sign of kidney dysfunction)
- unusual bleeding from the vagina
- swelling (oedema), chills and rigors

Rare side effects (may affect up to 1 in 1,000 people) include:

- angioedema (swelling mainly of the face, lip, tongue or throat, itching and rashes)

Some of these side effects are more common when capecitabine is used with other medicines for the treatment of cancer. Other side-effects seen in this setting are the following:

Common side-effects (may affect up to 1 in 10 people) include:

- decrease in blood sodium, magnesium or calcium, increase in blood sugar
- nerve pain
- ringing or buzzing in the ears (tinnitus), loss of hearing
- vein inflammation
- hiccups, change in voice
- pain or altered/abnormal sensation in the mouth, pain in the jaw
- sweating, night sweats
- muscle spasm
- difficulty in urination, blood or protein in the urine
- bruising or reaction at the injection site (caused by medicines given by injection at the same time)

Rare side effects (may affect up to 1 in 1,000 people) include:

- narrowing or blockage of tear duct (lacrimal duct stenosis)
- liver failure
- inflammation leading to dysfunction or obstruction in bile secretion (cholestatic hepatitis)
- specific changes in the electrocardiogram (QT prolongation)
- certain types of arrhythmia (including ventricular fibrillation, torsade de pointes, and bradycardia)
- eye inflammation causing eye pain and possibly eyesight problems
- inflammation of the skin causing red scaly patches due to an immune system illness

Very rare side effects (may affect up to 1 in 10,000 people) include:

- severe skin reaction such as skin rash, ulceration and blistering which may involve ulcers of the mouth, nose, genitalia, hands, feet and eyes (red and swollen eyes)

Reporting of side effects

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Ecansya

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

For Aluminium-aluminium blisters

This medicine does not require any special storage conditions.

For PVC/PVdC-aluminium blisters

Do not store above 30°C.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Ecansya contains

- The active substance is capecitabine. Each film-coated tablet contains 150 mg, 300 mg or 500 mg capecitabine.
- The other ingredients (excipients) are:

Tablet core

Lactose, croscarmellose sodium, hypromellose (E-5), microcrystalline cellulose, magnesium stearate.

Film-coating

Ecansya 150 mg film-coated tablets

Hypromellose (6cps), titanium dioxide (E171), yellow iron oxide, red iron oxide (E172), talc.

Ecansya 300 mg film-coated tablets

Hypromellose (6cps), titanium dioxide (E171), talc.

Ecansya 500 mg film-coated tablets

Hypromellose (6cps), titanium dioxide (E171), yellow iron oxide, red iron oxide (E172), talc.

See section 2 "Ecansya contains lactose".

What Ecansya looks like and contents of the pack

Ecansya 150 mg film-coated tablets (tablets) are light peach coloured, oblong shaped, biconvex tablets of 11.4 mm in length and 5.3 mm in width, debossed with "150" on one side and plain on other side.

Ecansya 300 mg film-coated tablets (tablets) are white to off white, oblong shaped, biconvex tablets of 14.6 mm in length and 6.7 mm in width, debossed with "300" on one side and plain on other side.

Ecansya 500 mg film-coated tablets (tablets) are peach coloured, oblong shaped, biconvex tablets of 15.9 mm in length and 8.4 mm in width, debossed with "500" on one side and plain on other side.

Ecansya is available in blisters (Aluminium-aluminium or PVC/PVdC-aluminium) containing 30, 60 or 120 film-coated tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder

KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

Manufacturer

Pharmacare Premium Limited, HHF 003, Hal Far Industrial Estate Birzebbugia, BBG 3000, Malta
KRKA, d.d., Novo mesto, Šmarješka cesta 6, 8501 Novo mesto, Slovenia

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

België/Belgique/Belgien

KRKA Belgium, SA.

Tél/Tel: + 32 (0) 487 50 73 62

Lietuva

UAB KRKA Lietuva

Tel: + 370 5 236 27 40

България

КРКА България ЕООД

Тел.: + 359 (02) 962 34 50

Luxembourg/Luxemburg

KRKA Belgium, SA.

Tél/Tel: + 32 (0) 487 50 73 62 (BE)

Česká republika

KRKA ČR, s.r.o.

Tel: + 420 (0) 221 115 150

Magyarország

KRKA Magyarország Kereskedelmi Kft.

Tel.: + 36 (1) 355 8490

Danmark

KRKA Sverige AB

Tlf: + 46 (0)8 643 67 66 (SE)

Malta

E. J. Busuttil Ltd.

Tel: + 356 21 445 885

Deutschland

Nederland

TAD Pharma GmbH
Tel: + 49 (0) 4721 606-0

Eesti

KRKA, d.d., Novo mesto Eesti filiaal
Tel: + 372 (0) 6 671 658

Ελλάδα

QUALIA PHARMA S.A.
Τηλ: + 30 210 6256177

España

KRKA Farmacéutica, S.L.
Tel: + 34 911 61 03 81

France

KRKA France Eurl
Tél: + 33 (0)1 57 40 82 25

Hrvatska

KRKA - FARMA d.o.o.
Tel: + 385 1 6312 100

Ireland

KRKA Pharma Dublin, Ltd.
Tel: + 353 1 293 91 80

Ísland

LYFIS ehf.
Sími: + 354 534 3500

Italia

KRKA Farmaceutici Milano S.r.l.
Tel: + 39 02 3300 8841

Κύπρος

KI.PA. (PHARMACAL) LIMITED
Τηλ: + 357 24 651 882

Latvija

KRKA Latvija SIA
Tel: + 371 6 733 86 10

KRKA Belgium, SA.
Tel: + 32 (0) 487 50 73 62 (BE)

Norge

KRKA Sverige AB
Tlf: + 46 (0)8 643 67 66 (SE)

Österreich

KRKA Pharma GmbH, Wien
Tel: + 43 (0)1 66 24 300

Polska

KRKA-POLSKA Sp. z o.o.
Tel.: + 48 (0)22 573 7500

Portugal

KRKA Farmacêutica, Sociedade Unipessoal Lda.
Tel: + 351 (0)21 46 43 650

România

KRKA Romania S.R.L., Bucharest
Tel: + 4 021 310 66 05

Slovenija

KRKA, d.d., Novo mesto
Tel: + 386 (0) 1 47 51 100

Slovenská republika

KRKA Slovensko, s.r.o.
Tel: + 421 (0) 2 571 04 501

Suomi/Finland

KRKA Finland Oy
Puh/Tel: + 358 20 754 5330

Sverige

KRKA Sverige AB
Tel: + 46 (0)8 643 67 66 (SE)

United Kingdom (Northern Ireland)

Consilient Health Limited
Tel: + 353 (0)1 2057760

This leaflet was last revised in

Detailed information on this medicine is available on the European Medicines Agency web site:
<http://www.ema.europa.eu>.